

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
11 November 2004 (11.11.2004)

PCT

(10) International Publication Number  
**WO 2004/096148 A2**

(51) International Patent Classification<sup>7</sup>: **A61K**

(21) International Application Number:  
PCT/US2004/013099

(22) International Filing Date: 26 April 2004 (26.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/465,891 25 April 2003 (25.04.2003) US  
10/676,694 30 September 2003 (30.09.2003) US

(71) Applicant (for all designated States except US): **THE KENNETH S. WARREN INSTITUTE, INC.** [US/US]; 712 Kitchawan Road, Ossining, NY 10562 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRINES, Michael** [US/US]; 1 Wepawaug Road, Woodbridge, CT 06525 (US). **CERAMI, Anthony** [US/US]; 49 Bramblebush Drive, Croton-on-Hudson, NY 10520 (US). **GHEZZI, Pietro** [IT/IT]; Via Mac Mahon No. 82, I-20155 Milan (IT). **FIORDALISO, Fabio** [IT/IT]; Via Lorenzo di Credi 10, I-20149 Milan (IT). **FRATELLI, Maddalena** [IT/IT];

Via Costanza 4, I-20146 Milan (IT). **LEIST, Marcel** [DK/DK]; Kloverbladsgade 14.1, DK-2500 Valby (DK). **NIELSEN, Mette** [DK/DK]; Gladsaxevej 73.3. th, DK-2860 Soborg (DK). **SAGER, Thomas** [DK/DK]; Stormosevej 35, DK-2765 Smorum (DK). **PEDERSEN, Lars, Ostergaard** [DK/DK]; Ved Vigen 14, DK-2400 Copenhagen (DK). **GERWIEN, Jens** [DK/DK]; Haraldsgade 52, 1.tv, DK-2100 Copenhagen (DK). **COLEMAN, Thomas** [US/US]; 20 Emery Street, Mt. Kisco, NY 10549 (US).


(74) Agents: **CORUZZI, Laura, A.** et al.; Jones Day, 222 East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: TISSUE PROTECTIVE CYTOKINE RECEPTOR COMPLEX, ASSAYS FOR IDENTIFYING TISSUE PROTECTIVE COMPOUNDS AND USES THEREOF



(57) Abstract: The present invention is directed methods for identifying compounds that have a tissue protective activity using a heteromultimer receptor complex that mediates the tissue protective activities. The complex consists of at least one EPO-R in complex with at least one  $\beta$ c Receptor. These compounds used in the assays to identify tissue protective compounds include, but are not limited to, small molecules and biologics. The compounds identified using these assays can be used to treat or prevent various diseases, disorders, or conditions of the central and peripheral nervous systems as well as those of other erythropoietin-responsive or excitable cells, tissues, and organs.

WO 2004/096148 A2



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

- *without international search report and to be republished upon receipt of that report*

**TISSUE PROTECTIVE CYTOKINE RECEPTOR COMPLEX, ASSAYS FOR  
IDENTIFYING TISSUE PROTECTIVE COMPOUNDS AND USES THEREOF**

This application claims benefit of priority of copending U.S. Patent Application  
5 No. 10/676,694, filed September 30, 2003, which claims benefit of priority under 35  
U.S.C. § 119(e) to U.S. provisional patent Application No. 60/465,891, filed April 25,  
2003, the entire contents of each of which is incorporated herein by reference in their  
entirety.

**1. INTRODUCTION**

10 The present invention is directed to methods that target tissue protective  
cytokine receptor complexes having at least one beta common ( $\beta_c$ ) receptor (also known  
as CD131) and at least one erythropoietin (EPO) receptor for drug screening/discovery  
and methods of treatment or prophylaxis. In particular, the present invention is drawn to  
methods for identifying and screening for compounds that modulate the interaction of a  
15 tissue protective cytokine receptor complex and a tissue protective cytokine receptor  
complex ligand. The methods of the invention also encompass methods for treating or  
preventing a disease or disorder using compounds that modulate the activity of a tissue  
protective cytokine receptor complex such as those compounds identified by the  
screening methods of the invention. The invention also encompasses methods for  
20 enhancing excitable tissue function using compounds that modulate the activity of a  
tissue protective cytokine receptor complex such as those compounds identified by the  
screening methods of the invention.

## 2. BACKGROUND OF THE INVENTION

Several lines of evidence suggest that erythropoietin, as a member of the cytokine superfamily, performs important physiologic functions which are mediated through interaction with the erythropoietin receptor (EPO-R). These actions include  
5 production of red blood cells, mitogenesis, modulation of calcium influx into smooth muscles and neural cells, and effects on intermediary metabolism.

EPO-R is a 68kDa protein, and is part of the Type-1 cytokine receptor family. This family includes receptors for interleukin (IL)-IL2, IL3, IL4, IL5, IL6, IL7, IL9, IL11, granulocyte macrophage – colony stimulating factor (GM-CSF), granulocyte  
10 colony stimulating factor (G-CSF), Leukaemia Inhibiting Factor (LIF), Ciliary Neurotrophic Factor (CNTF), Thrombopoietin, Growth Hormone and Prolactin. These receptors are grouped together because of the homology of their extracellular domains. The conserved extracellular domain of these receptors has a length of approximately 200 amino acids, which contains four positionally conserved cysteine residues in the amino-  
15 terminal region (Cys 294, Cys 283, Cys 248, Cys 238) and a Trp-Ser-X-Trp-Ser motif located proximal to the transmembrane domain. The four cysteines appear to be critical to the maintenance and the structural integrity of the receptors (Murray, 1996, Harpers Biochemistry 24<sup>th</sup> ed. pp.524-526, Appilion & Lange, Ltd.; Caravella *et al.*, 1996, Protein: Struct. Funct. Gen. 24:394-401; ).

20 Like many of the receptors within the Type-1 cytokine receptor family, the EPO-R appears to be activated by interaction its endogenous ligand EPO. The first EPO-R in the dimer binds to EPO with a high affinity and the second EPO-R then binds to the complex with a low affinity. This dimerization of the EPO-R puts the Jak2 tyrosine kinases associated with EPO-R in close association, inducing their  
25 transphosphorylation. This activation leads to the tyrosine phosphorylation of several



proteins that subsequently activate several different pathways, such as phosphatidylinositol (PI) 3-kinase pathway, the Ras/MAP kinase pathway, and the STAT pathway. These pathways trigger the physiological functions mediated by erythropoietin (Kirito *et al.*, 2002, Blood 99:102-110; Livnah *et al.*, 1999, Science 283:987-990; 5 Naranda *et al.*, 2002, Endocrinology 143:2293-2302; Remy *et al.*, 1999, Science 283:990-993; and Yoshimura *et al.*, 1996, The Oncol. 1:337-339).

In addition to forming multimers, many of the members of the type-1 family incorporate one of three different signal transducing receptor components – gp130, beta common ( $\beta_c$  receptor), or the gamma subunit of the IL2 receptor ( $\gamma_c$  10 receptor) – into their receptor complexes. For example, the receptor complex for GM-CSF consists of GM-CSF receptor, two  $\beta_c$  receptors, and the GM-CSF ligand. The EPO-R has been known to form a complex with the  $\beta_c$  receptor (*See* Yutaka *et al.*, 1995, Bioch. Biophys. Res. Com. 208:1060-1066; Jubinsky *et al.*, 1997, Blood 90:1867-1873; D’Andrea *et al.*, 1998, J. Clin. Invest. 102:1951-1960). However, it has been reported 15 that these complexes failed to result in any physiologically relevant effect (Scott *et al.*, 2000, Blood, 96:1588-1590).

### 3. SUMMARY

The invention relates to targeting “tissue protective cytokine receptor complexes” (defined *infra*) for drug screening/discovery, and methods for achieving 20 tissue protection or repair in vivo, *e.g.*, for the prophylaxis or treatment of a variety of diseases, disorders and conditions, or for the improvement of function of excitable tissue in human subjects or other mammals. The invention is based, in part, on the applicants’ discovery that EPO receptor/ $\beta_c$  receptor complexes play a role in the protection of excitable tissues. Experiments described herein demonstrate that EPO had a protective

effect, *i.e.*, mitigated apoptosis, in wild-type cells that express both EPO receptors and  $\beta_c$  receptors; however, EPO had no protective effect in cells lacking the  $\beta_c$  receptors. In animal studies, ischemic spinal injury was inflicted in wild-type mice (which express both receptors) and in  $\beta_c$  receptor knock-out mice (which do not express the  $\beta_c$  receptor).

5 The administration of carbamylated EPO (which is non-erythropoietic and does not bind EPO receptor dimers) resulted in an improved recovery in the wild-type mice. However, neither EPO nor carbamylated EPO had an effect on recovery in the  $\beta_c$  receptor knock-out mice. These results demonstrate that expression of the  $\beta_c$  receptor is essential to achieve a tissue protective effect.

10 In one aspect, the present invention is directed to the use of tissue protective cytokine receptor complexes, comprising an EPO receptor and a  $\beta_c$ , in cell-free screening assays to identify compounds that bind tissue protective cytokine receptor complexes. In other aspects, the present invention is directed to the use of tissue protective cytokine receptor complexes, comprising an EPO receptor and a  $\beta_c$ , in cell-  
15 based screening assays to identify compounds that modulate, *i.e.*, enhance or inhibit, the activity of tissue protective cytokine receptor complexes or exhibit a tissue protective activity. In other aspects, the present invention is directed to the use of tissue protective cytokine receptor complexes, comprising an EPO receptor and a  $\beta_c$ , in animal-based screening assays to identify compounds that exhibit a tissue protective activity dependent  
20 on tissue protective cytokine receptor complexes. In another particular aspect, the compounds identified by the screening methods of the invention modulate the interaction of a tissue protective cytokine receptor complex and a ligand thereof and exhibit a tissue protective activity. In other embodiments, the invention provides for methods for treating or preventing a disease, disorder or condition comprising administering a  
25 compound that modulates the activity of a tissue protective cytokine receptor complex.

In another aspect, the invention provides for methods for enhancing the function of excitable cells, tissues, or organs by administering a compound that modulates the activity of a tissue protective cytokine receptor complex.

In certain embodiments of the screening methods of the invention, the test compound is contacted with a tissue protective cytokine receptor complex in the presence and/or in the absence of a ligand of the tissue protective cytokine receptor complex, such as, but not limited to EPO. Such assays can be used to identify compounds that modulate the interaction of a ligand and a tissue protective cytokine receptor complex.

Cell-free assays can be used to identify compounds that bind a tissue protective cytokine receptor complex. In such embodiments, tissue protective cytokine receptor complexes or a membrane preparation of such complexes is contacted with a test compound and an indication of the binding is measured. The assay may be a direct binding assay where a compound is contacted to a tissue protective cytokine receptor complex. In competitive binding assays two compounds (e.g., a test compound and a ligand, or two test compounds) are contacted to a tissue protective cytokine receptor complex to determine which of the two compounds has greater binding affinity. The assay may be a displacement binding assay where a first compound is contacted to a tissue protective cytokine receptor complex that binds to the complex and a second compound is added to the assay to determine if the second compound has the ability to displace the first bound compound. In binding assays, the binding can be measured by labeling the compounds added to the assay and/or labeling a tissue protective cytokine receptor complex.

Cell-based assays can be used to identify compounds that target a tissue protective cytokine receptor complex in the presence and/or absence of a ligand for a

tissue protective cytokine receptor complex. In such embodiments, a cell that expresses a tissue protective cytokine receptor complex is contacted with a test compound. To determine whether the test compound modulates the activity of a tissue protective cytokine receptor complex in the cell, a readout is observed or measured. The readout can be a simple measure of the binding of the test compound to the tissue protective cytokine receptor complex. In certain embodiments, the test compound is labeled and binding of the labeled test compound to the tissue protective cytokine receptor complex is measured by detecting the label attached to the test compound. In other embodiments, the readout is a phenotypic change such as a change in cell size, shape, or apoptosis of the cell. In yet other embodiments, the readout is expression of a reporter gene or a change in an indicator dye loaded into the cell or included in the culture media. In such embodiments, the cells employed in the assay are genetically engineered to express a reporter gene product upon activation of a tissue protective cytokine receptor complex. In other embodiments, the readout is determined by enzymatic reaction of the cell. In such embodiments, cells used in the assay after being contacted with a test compound are lysed. The cell lysates are then examined for indicators of the activation of a tissue protective cytokine receptor complex. For example, a cell lysates can be exposed to an antibody that binds the complex, which antibody can also attach to a support anchoring the complex. Other antibodies can then be added to the assay to detect whether the tissue protective cytokine receptor complex has been activated. For example, anti-phosphotyrosine antibodies can bind to phosphorylated portions of the complex and/or to downstream components of the EPO/ $\beta$  common receptor complex. In certain embodiments, antibodies that can bind to phosphorylated portions of the  $\beta$  common receptor portion of the complex and/or antibodies can bind to phosphorylated portions of the EPO receptor are used. In certain embodiments, cells expressing tissue protective

cytokine receptor complexes are contacted with antibodies that can bind to phosphorylated portions of the  $\beta$  common receptor and/or the EPO receptor. In certain embodiments, isolated tissue protective cytokine receptor complexes are contacted with antibodies that can bind to phosphorylated portions of the  $\beta$  common receptor and/or the EPO receptor. In certain embodiments, antibodies that can bind to phosphorylated portions of the EPO receptor are PY-JAK2 antibodies. As downstream cytokine activation targets are identified for tissue protective cytokine receptor complexes, the readout of the screening methods of the invention can be detection of phosphorylation of downstream cellular components.

10           The presence of downstream cellular products can be used to identify compounds that target a tissue protective cytokine receptor complex. For example, following contacting of a compound with a cell expressing a tissue protective cytokine receptor complex, the cell can be examined for the presence of other cellular products such as nucleolin or neuroglobin that are indicative of activation of a tissue protective  
15 cytokine receptor complex.

          In certain embodiments of the screening methods of the invention, the cell is recombinantly engineered to express at least one EPO receptor or  $\beta$  common receptor. In certain embodiments, the cell is genetically engineered to overexpress a nucleic acid comprising a nucleotide sequence that encodes a  $\beta$  common receptor polypeptide or a  
20 EPO receptor polypeptide. In certain embodiments, the cell is genetically engineered to overexpress a nucleic acid comprising a nucleotide sequence that encodes a  $\beta$  common receptor polypeptide and a nucleic acid comprising a nucleotide sequence that encodes a EPO receptor polypeptide. The overexpression may be achieved by operably linking the nucleic acids to a promoter genetic engineering may occur by standard transformation  
25 techniques, targeted homologous recombination to achieve gene activation of

endogenous genes, or random integration and selection of cells with the activated gene. In certain embodiments, the nucleotide sequence is derived from the same species as the cell. In certain embodiments, the cell endogenously expresses a  $\beta$  common receptor and/or a EPO receptor.

5                   In one embodiment, the invention provides for a method for identifying a compound that modulates the activity of a tissue protective cytokine receptor complex in a  $\beta$  common receptor knock-out cell, comprising administering the compound to a first cell, immediately followed by subjecting the cell to a harmful agent, wherein the first cell endogenously expresses a tissue protective cytokine receptor complex and  
10       administering the compound to a second cell immediately followed by subjecting the cell to the same harmful agent as the first cell, wherein the second cell is of the same type as the first cell but is deficient in expression of a tissue protective cytokine receptor complex (*i.e.*, lacking expression of  $\beta$  common receptor), such that if survival of the cell differs in the first cell as compared to the second cell, then the compound is one that  
15       targets a tissue protective cytokine receptor complex. In one embodiment, the cells are derived from a  $\beta$  common receptor knock-out animal.

                  Assays can be conducted in animal models to identify compounds that target a tissue protective cytokine receptor complex. In one embodiment, the invention provides for a method for identifying a compound that modulates a tissue protective  
20       activity in a mammal, comprising administering the compound to a first animal immediately following infliction of an injury, wherein the first animal endogenously expresses a tissue protective cytokine receptor complex and administering the compound to a second animal immediately following infliction of the same type of injury as inflicted to the first animal, wherein the second animal is deficient in expression of a  
25       tissue protective cytokine receptor complex, such that if recovery from the injury differs

in the first animal as compared to the second animal, then a compound that modulates a tissue protective activity is identified.

Cells expressing a tissue protective cytokine receptor complex that can be used in the screening methods of the invention include, but are not limited to, mammal, insect, eukaryotic, prokaryotic, fungi, or bacteria cells. In certain embodiments, human cells are used, such as, but not limited to, pre-B cells, 293 cells, 293T cells, HeLa cells, HepG2 cells, K562 cells, or 3T3 cells. In other embodiments mouse cells are used, such as, but not limited to, BaF3 cells. In other embodiments fungi cells are used, such as, but not limited to, cells from the genera *Saccharomyces*, *Schizosaccharomyces*, *Pichia*, and *Hansenula*.

According to another aspect of the invention, a method of treating or preventing a disease or disorder in a human is provided which comprises administering a therapeutically effective amount of a compound that modulates the activity of a tissue protective cytokine receptor complex to a human in need of such treatment or prevention. In certain embodiments, the disease or disorder is a disease or disorder of or affecting excitable tissue.

In one embodiment, the invention provides for a method for enhancing the function of excitable cells, tissues, or organs in a human comprising administering a therapeutically effective amount of a compound that modulates the activity of a tissue protective cytokine receptor complex to a human in need thereof, wherein said enhancing the function of excitable cells, tissues, or organs results in the enhancement of associative learning or memory.

In another embodiment, the invention provides for a method for modulating the activity of a tissue protective cytokine receptor complex in a human,

comprising administering a compound that modulates the activity of said complex to a human in need of such treatment or prevention.

In yet another embodiment, the invention provides for a method for treating or preventing a disease or disorder in a human comprising modulating a tissue protective cytokine receptor complex in a human.

In still another embodiment, the invention provides for a method for enhancing the function of excitable cells, tissues, or organs in a human comprising modulating a tissue protective cytokine receptor complex in a human.

In embodiments of the invention for treating or preventing a disease or disorder, enhancing the function of excitable cells, tissues, or organs, or modulating the activity of a tissue protective cytokine receptor complex the compound administered can be identified by the method of any one of the screening methods described herein above for identifying compounds that have a tissue protective effect or modulate the activity of a tissue protective cytokine receptor complex. In related embodiments, the compound is an antibody specific for the tissue protective cytokine receptor complex. In related embodiments, compound is an antibody specific for a tissue protective cytokine receptor complex ligand. In related embodiments, the compound is a small molecule, peptide, or a member of a library. In related embodiments, the compound binds to the tissue protective cytokine receptor complex. In related embodiments, the compound enhances the activity of the tissue protective cytokine receptor complex. In related embodiments, compound interferes with the activity of the tissue protective cytokine receptor complex. In related embodiments, the compound is administered in conjunction with an EPO. In certain embodiments, the compound administered is an EPO. In certain embodiments, the compound administered is not EPO. In related embodiments, the compound administered is not a modified EPO or an EPO mutant. Examples of modified EPO and



mutant EPO compounds include, but are not limited to, those described in section 5.4.1.1. In certain embodiments of the invention, where the compound administered is an EPO, a modified EPO, or an EPO derivative, the compound administered does not bind the EPO receptor. In related embodiments of the invention, where the compound administered is an EPO, a modified EPO, or an EPO derivative, the compound administered does not bind an EPO receptor dimer. In certain embodiments of the methods of the invention for treatment, prevention, enhancement of function of excitable tissue, and modulation of a tissue protective cytokine receptor complex, the compound administered is not an EPO, *i.e.*, any EPO, modified EPO, mutated EPO, or derivative thereof.

In certain other embodiments of the methods of the invention for treatment or prevention of a disease disorder, or condition, enhancement of function of excitable tissue, or modulation of a tissue protective cytokine receptor complex, the compound administered does not bind the EPO receptor. In certain other embodiments of the methods of the invention for treatment or prevention of a disease disorder, or condition, enhancement of function of excitable tissue, or modulation of a tissue protective cytokine receptor complex, the compound administered does not bind a dimer consisting of two EPO receptors. In certain other embodiments of the methods of the invention for treatment or prevention of a disease disorder, or condition, enhancement of function of excitable tissue, or modulation of a tissue protective cytokine receptor complex, the compound administered binds a tissue protective cytokine receptor complex.

In embodiments of the invention for treating or preventing a disease or disorder, the disease or disorder is a mood disorder, an anxiety disorder, depression, autism, an attention deficit hyperactivity disorder, Alzheimer's disease, aging, or

cognitive dysfunction. In related embodiments, the disease or disorder is caused by hypoxia, seizure disorders, neurodegenerative diseases, neurotoxin poisoning, multiple sclerosis, hypotension, cardiac arrest, radiation, or hypoglycemia. In related  
embodiments, the disease or disorder is ischemia or stroke. In related embodiments, the  
5 disease or disorder is a neurodegenerative condition. In related embodiments, the  
neurodegenerative condition is stroke, Alzheimer's disease, Parkinson's disease, cerebral  
palsy, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive  
function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism,  
retinal ischemia, aging, glaucoma or neuronal loss. In other related embodiments, the  
10 disease or disorder is a neuromuscular or muscular condition. In certain embodiments,  
the neuromuscular or muscular condition is of a heart tissue. In certain embodiments of  
the methods of the invention for treatment or prevention of a disease disorder, or  
condition, enhancement of function of excitable tissue, or modulation of a tissue  
protective cytokine receptor complex, the disease disorder, condition, or excitable cells,  
15 tissues, or organs, are not cerebral. In certain embodiments of the methods of the  
invention for treatment or prevention of a disease disorder, or condition, enhancement of  
function of excitable tissue, or modulation of a tissue protective cytokine receptor  
complex, the disease disorder, condition, or excitable cells, tissues, or organs, are not of  
the central nervous system including, but not limited to, renal tissues, cardiac tissues,  
20 endothelial tissues, retinal tissues, vascular tissues, lung tissues, pancreas tissues, bone  
tissues, skin tissues, embryonic tissues, fetal tissues, kidney tissues, striated muscle  
tissues, liver tissues, gastrointestinal tissues reproductive tract tissues, or endocrine  
tissues.

In embodiments of the invention for enhancing the function of excitable  
25 cells, tissues, or organs, the excitable tissue is central nervous system tissue or peripheral

nervous system tissue. In certain embodiments, the peripheral nervous system tissue is placental tissue, retinal tissue, or cardiac tissue.

In embodiments of the invention for treating or preventing a disease or disorder, enhancing the function of excitable cells, tissues, or organs, or modulating the activity of a tissue protective cytokine receptor complex the administration is oral, topical, intracranial, intraluminal, inhalation, nasal, parenteral, intravenous, peripherally, intraarterial, subcutaneous, intramuscular, intraperitoneal, submucosal or intradermal administration. In related embodiments, the administration is prior to a medical or surgical procedure. In certain embodiments, the surgical procedure is tumor resection, aneurysm repair, or a coronary artery bypass procedure. In certain embodiments of the invention, the medical procedure is labor or delivery.

### 3.1 TERMINOLOGY

As used herein, the term “tissue protective activity” refers to the effect of inhibiting or delaying damage or death of a cell, tissue, or organ. The tissue protective activity can be against various conditions, diseases, and cellular, organ, and/or tissue damage, for example, those described in section 5.5 . Tissue protective activity may be specific to excitable tissue, cells, and/or organs having a tissue protective cytokine receptor complex, such as, but not limited to, the tissues of the central nervous system.

The term “tissue protective cytokine receptor complex” as used herein means a complex comprising at least one erythropoietin receptor and at least one beta common receptor. The tissue protective cytokine receptor complex may contain multiple erythropoietin receptors and/or beta common receptors, as well as other types of receptors as described herein in section 5.1.3

The term “tissue protective cytokine receptor complex ligand” means any compound that binds a tissue protective cytokine receptor complex and activates the

complex or inhibits activation by a known activating ligand. A “tissue protective cytokine receptor complex ligand” can be any type of compound, such as, but not limited to, proteins, peptides, small molecules, organic molecules, or non-organic molecules. Non-limiting examples of tissue protective cytokine receptor complex ligands include

5 EPO, including mutants, chemical modifications, or tissue protective cytokine receptor complex-binding fragments thereof. Another example of a tissue protective cytokine receptor complex ligand is an antibody specific for the complex.

The term “preventing a disease, disorder, or condition” means delaying the onset, hindering the progress, hindering the appearance, protection against, inhibiting

10 or eliminating the emergence, or reducing the incidence, of such disease, disorder, or condition. Use of the term “prevention” is not meant to imply that all patients in a patient population administered a preventative therapy will never develop the disease, disorder, or condition targeted for prevention, but rather that the patient population will exhibit a reduction in the incidence of the disease, disorder, or condition. For example,

15 many flu vaccines are not 100% effective at preventing flu in those who administered the vaccine. Prevention also encompasses protection of excitable tissues. One skilled in the art can readily identify patients and situations for whom preventative therapy would be beneficial, such as, but not limited to, patients for whom surgery is planned, patients at risk for inherited diseases, disorders, or conditions, patients at risk for diseases,

20 disorders, or conditions precipitated by environmental factors, or portions of the population at risk for particular diseases, disorders, or conditions such as the elderly, infants, or those with weakened immune systems, or those patients with genetic or other risk factors for a disease, disorder, or condition.

The term “administered in conjunction with” in the context of the methods of the invention means administering a compound prior to, at the same time as, and/or subsequent to the onset of a disease, disorder, or condition.

#### 4. BRIEF DESCRIPTION OF THE FIGURES

5           **Figure 1** depicts the results of a competitive assay between EPO and carbamylated EPO that demonstrate that carbamylated EPO does not bind to the EPO-R.

**Figure 2** shows the presence of the  $\beta_c$  receptor within rat brain membranes.

**Figure 3** shows rat spinal cord tissue stained for the presence of the  $\beta_c$   
10    receptor.

**Figure 4** shows enhanced view of rat spinal cord tissue stained for the presence of the  $\beta_c$  receptor.

**Figure 5** shows the coprecipitation of the EPO-R and  $\beta_c$  receptor.

**Figure 6** shows the potential configuration of the tissue protective  
15    receptor complex.

**Figure 7** shows apoptosis in isolated cardiomyocytes from normal and  $\beta_c$  receptor knockout mice in the presence and absence of EPO.

**Figure 8** shows photographs of Western blotting of the cell proteins separated by polyacrylamide gels using phosphotyrosine (PY)-Jak2 antibodies. The top gel  
20    shows that phosphorylated Jak2 was present in cells stimulated with EPO at 5 nM and 50nM concentrations. The bottom gel shows a control performed where membranes were stripped and reprobed with an antibody against Jak2 to confirm equal loading.

**Figure 9** shows a graph of ligand binding affinity for BaF3 cells expressing the EPO receptor. Ligand concentration in nM (x-axis) is plotted against

binding (cpm) (y-axis) for BaF3 EPO receptor expressing cells contacted with EPO receptor ligands EPO and carbamylated EPO.

**Figure 10** shows a graph of ligand binding affinity for UT-7 cell membranes having the EPO receptor. Ligand concentration in nM (x-axis) is plotted against binding (cpm) (y-axis) for UT-7 EPO receptor expressing cell membranes contacted with EPO receptor ligands EPO and carbamylated EPO.

**Figure 11** shows histograms of, **11A**, hematocrit as measured by the percent volume of hematocrit (y-axis), and, **11B**, hemoglobin levels measured in concentration in mM (y-axis), in mice after administration of control (vehicle), EPO, carbamylated EPO, and asialoEPO (x-axis).

**Figure 12** shows a photograph of the SDS-PAGE gel with a 103 KD nucleolin protein circled.

**Figure 13** shows a graph of the BBB motor score (y-axis) versus time measured in days following spinal injury (x-axis) where five groups of mice, two wild type and three  $\beta_c$  knock-out, were administered either PBS, EPO, or carbamylated EPO immediately following spinal injury. The wild type group of mice administered carbamylated EPO exhibited increased motor scores throughout the 42 days post-injury time period in comparison to all other groups.

**Figure 14** shows a histogram of the clinical severity measured in area under curve (AUC) (y-axis) and groups of mice (x-axis) for the five groups described above in Figure 13. The wild type group of mice administered carbamylated EPO (column 1) exhibited a significant increase in clinical severity values (*i.e.*, less severity) in comparison to all other groups (columns 2-5).

## 5. DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to methods for use of a heteromultimer receptor complex that mediates the tissue protective activities of tissue protective compounds. The invention provides screening methods to identify such tissue protective compounds and use of such compounds in treatment or prevention of a disease, disorder, or condition, or use of such compounds in methods for enhancing the function of excitable cells, tissues, and organs. The invention also provides for methods of treatment or prevention of disease, disorders, or conditions by administering compounds that modulate the activity of a tissue protective cytokine receptor complex, such as, but not limited to, those identified by the screening methods of the invention. The invention also provides for methods of enhancing the function of excitable cells, tissues, or organs by administering compounds that modulate the activity of a tissue protective cytokine receptor complex, such as, but not limited to, those identified by the screening methods of the invention. The invention also provides for methods for modulating the activity of a tissue protective cytokine receptor complex by administering a compound that modulates or is targeted to the tissue protective cytokine receptor complex.

The invention is based on the discovery by the inventors that EPO receptor forms a complex with  $\beta_c$  receptor within excitable cells and tissues such as brain, spinal cord and heart. The complex consists of at least one EPO-R in a complex with at least one  $\beta_c$  receptor. In addition to the  $\beta_c$  receptor, other signal transducing receptors may be used including, but not limited to, the  $\gamma_c$  receptor and GP130, segments and portions of other Type-1 cytokine receptors, including, but not limited to, GM-CSF, IL-3, IL-5, *etc.*, and orphan receptors, including, but not limited to, ROR1, NR6, HM74, *etc.* The screening methods of the invention may use such tissue protective cytokine receptor complexes in assays for identifying compounds that modulate the activity of a

tissue protective cytokine receptor complex. Such assays include binding assays to identify compounds that bind to a tissue protective cytokine receptor complex or a ligand thereof. The discovery of the applicants provided the means to identify exceptionally effective compounds for use in treatment and prevention of disease, disorders, and conditions as well as enhancement of excitable cells, tissues, and organs.

Activation of a tissue protective cytokine receptor complex may lead to the upregulation of the production of protective proteins, including, but not limited to, nucleolin and globins such as neuroglobins and cytoglobin (histoglobins). In certain embodiments of the methods of the invention described herein above where tissue protective activity is assayed, the step of assaying the identified compound for tissue protective activity comprises detecting the presence of nucleolin in the cell.

In certain embodiments of the methods of the invention described herein where tissue protective activity is assayed, the step of assaying the identified compound for tissue protective activity comprises detecting or measuring an increased level of activity of neuroglobin or cytoglobin in a cell. Accordingly, the screening methods for identifying compounds with tissue protective activity may utilize detection of such upregulated proteins.

This tissue protective cytokine receptor complex and downstream regulators may be used in assays to identify tissue protective compounds, including, but not limited to, small molecules and biologics. The compounds identified using these assays can be used to treat various conditions of the central and peripheral nervous systems as well as those of other excitable cells, tissues, and organs.

## 5.1 THE TISSUE PROTECTIVE CYTOKINE RECEPTOR COMPLEX

The receptor complex for use in the methods of the invention comprises at least one EPO receptor and at least one  $\beta_c$  receptor. Various types of naturally-occurring,



modified, and/or mutant EPO receptors and  $\beta_c$  receptors may be combined to form such complexes of varying numbers of receptors. Non-limiting examples of tissue protective cytokine receptor complexes are described below.

### 5.1.1 EPO-R

5 Mammalian EPO-R may be used in the present invention. EPO-R cDNA has been isolated from mouse liver, Tojo *et al.*, 1987, Biochem. Biophys. Res. Comm. 148: 443-48 and from human fetal liver (Jones *et al.*, 1990, Blood 76:31-35; Winkelmann *et al.*, 1990, Blood 76:24-30). This human cDNA encodes a polypeptide chain of about 55 kDa MW and has about 508 amino acids. Other genomic clones of  
10 human EPO-R that have been isolated and sequenced are also contemplated by the present invention (Penny and Forget, 1991, Genomics 11:974-80; Noguchi *et al.*, 1991, Blood 78:2548-2556). The common structure of these EPO-R consist of about 24 amino acid residues in a signal peptide, about 226 amino acids in an extracellular domain, about 23 amino acids in a membrane-spanning domain, and about 235 amino acids in a  
15 cytoplasmic domain (D'Andrea and Zon, 1990, J. Clin. Invest., 86:681-687; Jones *et al.*, 1990, Blood, 76:31-35; Penny and Forget, 1991, Genomics, 11:974-80). The mature human EPO-R protein has about 484 amino acids. EPO-R is commercially available from BD Biosciences Clontech (Palo Alto, CA) and has gene bank accession number M60459 and SwissProt accession number P19235.

20 The forms of EPO-R useful in the practice of the present invention encompass naturally occurring, synthetic and recombinant forms of human and other mammalian EPO-R related molecules. In addition, EPO-R forms useful in the practice of the present invention include proteins that represent functionally equivalent gene products. Such a functionally equivalent gene products include mutant EPO-R and  
25 chemically modified EPO-R. Mutant EPO-R may contain deletions, including internal

deletions, additions, including additions yielding fusion proteins, or conservative substitutions of amino acid residues within and/or adjacent to the amino acid sequence, but that result in a “silent” change, in that the change produces a functionally equivalent EPO-R. Such amino acid substitutions may be made on the basis of similarity in

5 polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids including glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine,  
10 lysine, and histidine; and negatively charged (acidic) amino acids include aspartic and glutamic acid. Other known EPO-R mutations such as those disclosed in U.S. Patent No. 5,292,654 are encompassed by the present invention as well.

Further, soluble (truncated) forms of the EPO receptor containing only the extracellular domain are also contemplated for use in the present invention such as those  
15 disclosed in Harris *et al.* 1992, J. Biol. Chem. 267:15205; Yang & Jones, 1993, Blood 82:1713; and U.S. Patent Application Publication No. 20020031806.

### 5.1.2 $\beta_c$ Receptor

The  $\beta_c$  receptor is typically identified as a signaling receptor subunit associated with human GM-CSF, IL-3 and IL-5. In addition to these cytokines, the  $\beta_c$   
20 receptor (*e.g.*, Swiss accession no. P32927 or Genebank No. M59941) has been demonstrated to bind EPO-R in certain instances. The conserved region of the  $\beta_c$  receptor constitutes all or part of the extracellular ligand-binding region. The present invention is based, in part, on the demonstration by the Applicants that the EPO-R and the  $\beta_c$  receptor co-exist within several excitable cells, such as the brain (see Figure 2) and  
25 spinal cord (see Figures 3 and 4). Applicants also discovered that complexes of the

EPO-R and the  $\beta_c$  receptor play a role in a mechanism that lead to the physiologic process of protection and repair of tissues that express these complexes (see Figures 13 and 14).

The forms of  $\beta_c$  receptor useful in the practice of the present invention encompass naturally occurring, synthetic and recombinant forms of human and other mammalian  $\beta_c$  receptor related molecules. In addition,  $\beta_c$  receptor forms useful in the practice of the present invention include proteins that represent functionally equivalent gene products. Such functionally equivalent gene products may include mutant  $\beta_c$  receptor and chemically modified  $\beta_c$  receptor. The mutant  $\beta_c$  receptors may contain deletions, including internal deletions, additions, including additions yielding fusion proteins, or conservative substitutions of amino acid residues within and/or adjacent to the amino acid sequence, but that result in a "silent" change, resulting in a functionally equivalent  $\beta_c$  receptor. Mutated  $\beta_c$  receptors include, but are not limited to, I374N, F1A,  $\Delta$ GA, H544R, V449, A459D, L445Q, CRD4 point mutations (I374N, L356P, W358N, Q375P, Y376N, W383R, and L399P), and extracellular truncations (1GH7). (See R.J. D'Andrea and T.J. Gonda, 2000, *Experimental Hematology* 28:231-243; D'Andrea *et al.*, 1998, *J. Clin. Invest.* 102:1951-1960; Jenkins *et al.*, 1999, *J. Biol. Chem.* 274:8669-8677).

It is also contemplated that other known signaling receptor subunits including, but not limited to,  $\gamma_c$  receptor and GP-130, and orphan receptors may be used with the present invention as well.

### 5.1.3 Structure of the Heteromultimer Receptor Complex

The  $\beta_c$  receptor forms a multimer with many of the receptors it was previously known to bind with, *i.e.*, GM-CSF, IL-3, and IL-5. For example, the receptor complex for GM-CSF consists of GM-CSF receptor, two  $\beta_c$  receptors, and the GM-CSF

ligand. The EPO-R has been known to form a complex with the  $\beta_c$  Receptor. (See Yutaka *et al.*, 1995, Biochem. Biophys. Res. Com. 208:1060-1066; Jubinsky *et al.*, 1997, Blood, 90:1867-1873; D'Andrea *et al.*, 1998, J. Clin. Invest. 102:1951-1960.) Similarly, according to the present invention, the  $\beta_c$  receptor forms a heteromultimer with the EPO-R to form a tissue protective receptor complex. This is based on immunoprecipitation studies demonstrating that the  $\beta_c$  receptor and EPO-R receptors coprecipitate (see Figure 5).

This tissue protective receptor complex includes at least one EPO-R and at least one  $\beta_c$  receptor. Preferably, as depicted in Figure 6, the tissue protective receptor complex will include greater than one  $\beta_c$  receptor. Tissue protective receptor complex configurations are also contemplated by the present invention that include greater than one EPO-R. Examples of tissue protective receptor complexes contemplated by the present invention include, but are not limited to, EPO-R<sub>2</sub>/  $\beta_c$  receptor<sub>2</sub>, EPO-R/  $\beta_c$  receptor<sub>2</sub>, and EPO-R/  $\beta_c$  receptor<sub>3</sub>. The most suggested configuration of the tissue protective receptor complex is EPO-R/  $\beta_c$  receptor<sub>2</sub> as depicted in Figure 6. As shown therein, without being bound by any particular theory, a possible mechanism by which the tissue protective compound binds to the tissue protective receptor complex involves the tissue protective compound first binding with the EPO-R and then subsequently two  $\beta_c$  receptors binding to the complex. This depiction is merely illustrative of one method by which a ligand may bind to the tissue protective receptor complex and those of ordinary skill in the art could reasonably envision other mechanisms by which binding can occur – the receptor complex may be preformed (EPO-R and two  $\beta_c$  receptor) prior to binding the ligand, *etc.*

The receptor polypeptides of the present invention, including full-length receptors, receptor fragments (*e.g.* ligand-binding fragments), and fusion polypeptides

can be produced in genetically engineered host cells according to conventional techniques. For example, Erythropoietin binding fragments of the EPO receptor have been identified (Barone *et al.*, J. Biol. Chem. 272:4985-4992). Such fragments that also complex with a  $\beta_c$  receptor may be used in the screening methods of the invention.

## 5 5.2 SCREENING ASSAYS

Tissue protective cytokine receptor complexes, such as, but not limited to, those described above can be used in screening assays to identify compounds having a tissue protective activity. Compounds identified in these assays may modulate the activity of a complex, *e.g.*, bind to a tissue protective cytokine receptor complex, or  
10 modulate the interaction of a tissue protective cytokine receptor complex ligand and the complex. In addition, the compounds identified may themselves have a tissue protective activity.

This invention is particularly useful for screening compounds by using an EPO receptor- $\beta_c$  receptor complex in any of a variety of drug screening techniques. The  
15 receptor complex employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One method of drug screening utilizes eukaryotic cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed  
20 form, can be used for standard binding assays. One may measure, for example, the formation of complexes between receptor and the agent being tested or examine the diminution in complex formation between the tissue protective cytokine receptor complex and an appropriate cell line, which are well known in the art. For example, see examples sections 6 and 7 below.

The tissue protective cytokine receptor complexes described above in section 5.1 can similarly be used in assays to determine biological activity, including in a panel of multiple receptors for high-throughput screening, as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of a tissue protective compound in biological fluids, as markers for tissues in which the corresponding tissue protective compound is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state), and, of course, to isolate correlative ligands.

Methods for performing the assays listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

In certain embodiments, the assays of the invention described herein may be used to identify a compound that interacts with a tissue protective cytokine receptor complex. Such assays may be carried out in the absence of a tissue protective cytokine receptor complex ligand. The interaction may be, for example, binding of the test compound to tissue protective cytokine receptor complex or activity of a tissue protective cytokine receptor complex. In such embodiments, the test compound is contacted with a tissue protective cytokine receptor complex. The compound identified is one that produces a readout, such as, but not limited to, activation of the tissue protective cytokine receptor complex, transcription of a reporter gene, cell proliferation, or a tissue protective activity. Such assays are typically performed with a control, wherein the compound is not contacted to the tissue protective cytokine receptor

complex. Where there is a difference in the readout with and without the compound, a compound that interacts with the tissue protective cytokine receptor complex is identified. Alternatively, standard levels of readout, such as, but not limited to, activity of the tissue protective cytokine receptor complex, can be measured prior to contacting  
5 the tissue protective cytokine receptor complex with the test compound, and the difference in readout before and after addition of the compound can be used to identify compounds.

In certain other embodiments, the assays of the invention may be used to identify compounds that modulate, *i.e.*, enhance or interfere with, the interaction between  
10 a tissue protective cytokine receptor complex ligand and a tissue protective cytokine receptor complex. Such assays may be carried out in the presence of a tissue protective cytokine receptor complex ligand, such as EPO or an antibody specific to the tissue protective cytokine receptor complex. In such embodiments, the difference in the level of readout in the presence and absence of the test compound may be detected. The level  
15 of readout can then be compared to that in the absence of EPO. If the difference in the level of readout detected with and without the compound is dependent on the presence of EPO, then a compound that modulates the interaction of a tissue protective cytokine receptor complex with its ligand is identified. For example, if the difference in level of readout detected results from an increase in readout in the presence of a compound,  
20 wherein the increase is dependent on the presence of EPO, an agonist of the interaction between a tissue protective cytokine receptor complex and its ligand is identified. Alternatively, if the difference in level of readout detected results from a decrease in readout in the presence of a compound, wherein the decrease is dependent on the presence of EPO, then an antagonist of the interaction between a tissue protective  
25 cytokine receptor complex and its ligand is identified. The situation is reversed in

embodiments where the readout is a negative readout. For example, a readout may be present when a compound has no effect on the tissue protective cytokine receptor complex and is absent when a compound with such an effect is incorporated into the assay, then a negative readout may indicate that a compound with the desired effect has  
5 been identified.

Any or all of these assays that utilize a tissue protective cytokine receptor complex are capable of being developed into reagent grade or kit format for commercialization as research and/or clinical products.

### 5.2.1 Cell-Free Assays

10 The tissue protective cytokine receptor complexes described herein may be used in cell-free assays to identify compounds having a tissue protective activity. In certain embodiments of cell-free assays, the tissue protective cytokine receptor complexes may be free in solution or attached to a solid support. The activity of a tissue protective cytokine receptor complex may be measured in a variety of ways. For  
15 example, activity may be the compound's ability to bind a tissue protective cytokine receptor complex or the binding affinity of a tissue protective cytokine receptor complex and a compound. In certain embodiments, the tissue protective cytokine receptor complex is added to binding assays in the form of isolated membranes containing the tissue protective cytokine receptor complex.

20 In one embodiment, the invention provides for a method for identifying a compound that modulates a tissue protective activity comprising contacting a test compound with a tissue protective cytokine receptor complex, by measuring the level of tissue protective cytokine receptor complex activity, identifying a test compound which increases or decreases the level of tissue protective cytokine receptor complex activity as  
25 compared to the level of tissue protective cytokine receptor complex activity measured in



the absence of the test compound, and assaying the identified test compound for tissue protective activity.

In a direct binding assay, either the ligand and/or the tissue protective cytokine receptor complex is contacted with a test compound under conditions that allow binding of the test compound to the ligand or the receptor. The binding may take place in solution or on a solid surface. Preferably, the test compound is previously labeled for detection. Any detectable compound may be used for labeling, such as but not limited to, a luminescent, fluorescent, or radioactive isotope or group containing same, or a nonisotopic label, such as an enzyme or dye. After a period of incubation sufficient for binding to take place, the reaction is exposed to conditions and manipulations that remove excess or non-specifically bound test compound. Typically, it involves washing with an appropriate buffer. Finally, the presence of a ligand bound to the test compound (*e.g.*, EPO-test compound) or a the tissue protective cytokine receptor complex bound to the test compound is detected.

In a competition binding assay, test compounds are assayed for their ability to disrupt or enhance the binding of the ligand (*e.g.*, EPO) to the tissue protective cytokine receptor complex. Labeled ligand (*e.g.*, EPO or carbamylated EPO) may be mixed with the tissue protective cytokine receptor complex or a fragment thereof, and placed under conditions in which the interaction between them would normally occur, with and without the addition of the test compound. The amount of labeled ligand (*e.g.*, EPO) that binds the tissue protective cytokine receptor complex may be compared to the amount bound in the presence or absence of test compound.

In a preferred embodiment, to facilitate complex formation and detection, the binding assay is carried out with one or more components immobilized on a solid surface. In various embodiments, the solid support could be, but is not restricted to,

polycarbonate, polystyrene, polypropylene, polyethylene, glass, nitrocellulose, dextran, nylon, polyacrylamide and agarose. The support configuration can include beads, membranes, microparticles, the interior surface of a reaction vessel such as a microtiter plate, test tube or other reaction vessel. The immobilization of the tissue protective cytokine receptor complex, or other component, can be achieved through covalent or non-covalent attachments. In one embodiment, the attachment may be indirect, *i.e.* through an attached antibody. In another embodiment, the tissue protective cytokine receptor complex and negative controls are tagged with an epitope, such as glutathione S-transferase (GST) so that the attachment to the solid surface can be mediated by a commercially available antibody such as anti-GST (Santa Cruz Biotechnology).

For example, such an affinity binding assay may be performed using a tissue protective cytokine receptor complex which is immobilized to a solid support. Typically, the non-mobilized component of the binding reaction, in this case either ligand (*e.g.*, EPO or carbamylated EPO) or the test compound, is labeled to enable detection. A variety of labeling methods are available and may be used, such as luminescent, chromophore, fluorescent, or radioactive isotope or group containing same, and nonisotopic labels, such as enzymes or dyes. In a preferred embodiment, the test compound is labeled with a fluorophore such as fluorescein isothiocyanate (FITC, available from Sigma Chemicals, St. Louis).

The labeled test compounds, or ligand (*e.g.*, EPO or carbamylated EPO) plus test compounds, are then allowed to contact with the solid support, under conditions that allow specific binding to occur. After the binding reaction has taken place, unbound and non-specifically bound test compounds are separated by means of washing the surface. Attachment of the binding partner to the solid phase can be accomplished in various ways known to those skilled in the art, including but not limited to chemical

cross-linking, non-specific adhesion to a plastic surface, interaction with an antibody attached to the solid phase, interaction between a ligand attached to the binding partner (such as biotin) and a ligand-binding protein (such as avidin or streptavidin) attached to the solid phase, and so on.

5                    Preferably, the tissue protective cytokine receptor complex is added to binding assays in the form of intact cells that express the tissue protective cytokine receptor complex (see assays described in the sections 5.2.2 ), or isolated membranes containing the tissue protective cytokine receptor complex. Thus, direct binding to the tissue protective cytokine receptor complex or the ability of a test compound to modulate  
10 a ligand-tissue protective cytokine receptor complex (*e.g.*, EPO-tissue protective cytokine receptor complex) may be assayed in intact cells in culture or in animal models in the presence and absence of the test compound. A labeled ligand (*e.g.*, EPO or carbamylated EPO) may be mixed with crude extracts obtained from such cells, and the test compound may be added. Isolated membranes may be used to identify compounds  
15 that interact with the tissue protective cytokine receptor complex. For example, in a typical experiment using isolated membranes, cells may be genetically engineered to express the tissue protective cytokine receptor complex. Membranes can be harvested by standard techniques and used in an *in vitro* binding assay. Labeled ligand (*e.g.*, <sup>125</sup>I-labeled EPO) is bound to the membranes and assayed for specific activity; specific  
20 binding is determined by comparison with binding assays performed in the presence of excess unlabeled (cold) ligand.

                    The soluble tissue protective cytokine receptor complex ligand may also be recombinantly expressed and utilized in non-cell based assays to identify compounds that bind to the tissue protective cytokine receptor complex ligand. Alternatively, a  
25 ligand binding domain of the recombinantly expressed tissue protective cytokine

receptor complex, or a fragment of the tissue protective cytokine receptor complex, can be used in the non-cell based screening assays. In another alternative embodiment,

peptides corresponding to one or more of the binding domains of the tissue protective cytokine receptor complex, or fusion proteins containing one or more of the binding

5 domains of the tissue protective cytokine receptor complex can be used in non-cell based assay systems to identify compounds that bind to the cytoplasmic portion of the tissue protective cytokine receptor complex; such compounds may be useful to modulate a signal transduction pathway of the tissue protective cytokine receptor complex. In non-cell based assays the recombinantly expressed tissue protective cytokine receptor  
10 complex is attached to a solid substrate such as a test tube, microtiter well or a column, by means well known to those in the art (see Ausubel *et al.*, 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY). The test compounds are then assayed for their ability to bind to the tissue protective cytokine receptor complex.

Alternatively, the binding reaction may be carried out in solution. In this  
15 assay, the labeled component is allowed to interact with its binding partner(s) in solution. If the size differences between the labeled component and its binding partner(s) permit such a separation, the separation can be achieved by passing the products of the binding reaction through an ultrafilter whose pores allow passage of unbound labeled component but not of its binding partner(s) or of labeled component bound to its partner(s).  
20 Separation can also be achieved using any reagent capable of capturing a binding partner of the labeled component from solution, such as an antibody against the binding partner, a ligand-binding protein which can interact with a ligand previously attached to the binding partner, and so on.

In one embodiment, for example, a phage library can be screened by  
25 passing phage from a continuous phage display library through a column containing

purified tissue protective cytokine receptor complex, or a fragment, or domain, thereof, linked to a solid phase, such as plastic beads. By altering the stringency of the washing buffer, it is possible to enrich for phage that express peptides with high affinity for the tissue protective cytokine receptor complex. Phage isolated from the column can be  
5 cloned and the affinities of the short peptides can be measured directly. Sequences for more than one oligonucleotide can be combined to test for even higher affinity binding to the tissue protective cytokine receptor complex or its complex with its ligand. Knowing which amino acid sequences confer the strongest binding to the tissue protective cytokine receptor complex, computer models can be used to identify the molecular  
10 contacts between the tissue protective cytokine receptor complex and the test compound. This will allow the design of non-protein compounds which mimic those contacts. Such a compound may have the same activity of the peptide and can be used therapeutically, having the advantage of being efficient and less costly to produce.

In another specific embodiment of this aspect of the invention, the solid  
15 support is membranes containing the tissue protective cytokine receptor complex attached to a microtiter dish. Test compounds, for example, cells that express library members, are cultivated under conditions that allow expression of the library members in the microtiter dish. Library members that bind to the protein (or nucleic acid or derivative) are harvested. Such methods, are described by way of example in Parmley  
20 and Smith, 1988, *Gene* 73:305-318; Fowlkes *et al.*, 1992, *BioTechniques* 13:422-427; PCT Publication No. WO 94/18318; and in references cited hereinabove.

Finally, the label remaining on the solid surface may be detected by any detection method known in the art. For example, if the test compound is labeled with a fluorophore, a fluorimeter may be used to detect complexes.

Various *in vitro* assays can be used to identify and verify the ability of a compound to bind a tissue protective cytokine receptor complex, to modulate the interaction of a tissue protective cytokine receptor complex and a tissue protective cytokine receptor complex ligand, or modulate the activity of a tissue protective cytokine receptor complex.

In another embodiment of the present invention, interactions between the tissue protective cytokine receptor complex or ligand (*e.g.*, EPO or carbamylated EPO) and a test compound may be assayed *in vitro*. Known or unknown molecules are assayed for specific binding to the tissue protective cytokine receptor complex, or fragments thereof, under conditions conducive to binding, and then molecules that specifically bind to the tissue protective cytokine receptor complex are identified. The two components can be measured in a variety of ways. One approach is to label one of the components with an easily detectable label, place it together with a test component(s) under conditions that allow binding to occur, perform a separation step which separates bound labeled component from unbound labeled component, and then measure the amount of bound component. In one embodiment, the tissue protective cytokine receptor complex can be labeled and added to a test agent, using conditions that allow binding to occur. Binding of the test agent can be determined using polyacrylamide gel analysis to compare complexes formed in the presence and absence of the test agent.

Multiple *in vitro* assays can be performed simultaneously or sequentially to assess the effect of a compound on a tissue protective cytokine receptor complex-mediated process. The assessment can be made by measuring or detecting tissue protective cytokine receptor complex activity, for example, binding of a test compound to a tissue protective cytokine receptor complex, cell proliferation, cell differentiation, and upregulation of a protective protein, *e.g.*, globin. In a preferred embodiment, the *in*

*vitro* assays described herein are performed in a high throughput format (*e.g.*, in microtiter plates).

### 5.2.2 Cell-Based Assays

A tissue protective cytokine receptor complex may be expressed in a cultured cell, and the cell can then be used to screen for ligands for the receptor complex, including the natural ligand, as well as agonists and antagonists of the natural ligand.

To summarize the approach to the cell-based assay, a cDNA or gene encoding the receptor is combined with other genetic elements required for its expression (*e.g.*, a transcription promoter), and the resulting expression vector is inserted into a host cell. A vector encoding an EPO receptor or a  $\beta$  common receptor may be synthesized and used to transform or transfect a host cell with the vector of interest for use in the methods of the invention. The use of stable transformants is preferred.

Cells that express the DNA and produce functional receptor are selected and used with one of a variety of possible screening systems, as further disclosed below.

An example of a cell-based method for identifying a compound that modulates the interaction of a tissue protective cytokine receptor complex and a ligand thereof comprises the following steps: (a) contacting a test compound with the ligand and an tissue protective cytokine receptor complex-expressing cell; and (b) measuring the level of tissue protective cytokine receptor complex activity in the cell, such that if the level of activity measured in (b) differs from the level of tissue protective cytokine receptor complex activity in the absence of the test compound, then a compound that modulates the interaction of a tissue protective cytokine receptor complex and its ligand is identified.

Another example for a method for identifying a compound that modulates a tissue protective activity comprises contacting a test compound with a cell which is

recombinantly engineered to express a EPO receptor or a  $\beta$  common receptor, wherein said cell is transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a  $\beta$  common receptor or a EPO receptor polypeptide, measuring the level of tissue protective cytokine receptor complex activity

5 in the cell, identifying a test compound that increases or decreases the level of tissue protective cytokine receptor complex activity in the cell relative to the level of tissue protective cytokine receptor complex activity measured in a control cell, wherein the control cell is of the same cell type as the cell contacted with the test compound and is not transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably  
10 linked to a promoter, and (ii) encodes a  $\beta$  common receptor or a EPO receptor polypeptide, and assaying the identified test compound for a tissue protective activity.

Alternatively, cell-based assays may be used to identify compounds that interact with a tissue protective cytokine receptor complex. For example, a test compound can be contacted with a tissue protective cytokine receptor complex-

15 expressing cell and the activity of the tissue protective cytokine receptor complex, as measured by binding affinity, cell proliferation, or the presence of a reporter gene transcript, can be determined. If the activity of the tissue protective cytokine receptor complex differs, *e.g.*, is increased or decrease, in comparison to the activity of tissue protective cytokine receptor complex in the absence of the test compound, then a  
20 compound that interacts with a tissue protective cytokine receptor complex is identified.

In certain embodiments of the methods of the invention described herein above where tissue protective cytokine receptor complex activity is measured, such activity is measured by cell proliferation or cell differentiation.

In certain embodiments of the methods of the invention described herein  
25 above where tissue protective cytokine receptor complex activity is measured, such



activity measured is the ability of the tissue protective cytokine receptor complex to interact with a tissue protective cytokine receptor complex ligand.

Cell-based screening assays are useful for detecting both agonist and antagonist ligands. Agonists (including the natural ligand) and antagonists have enormous potential in both *in vitro* and *in vivo* applications. Compounds identified as receptor agonists are useful for stimulating proliferation and development of target cells *in vitro* and *in vivo*. For example, agonist compounds are useful as components of defined cell culture media, and may be used alone or in combination with other cytokines and hormones to replace serum that is commonly used in cell culture. Agonists may be useful in specifically promoting the growth and/or development of EPO-responsive cells (including, but not limited to, nervous, heart, and retinal derived cells) in culture.

Antagonists are useful as research reagents for characterizing sites of ligand-receptor interaction. *In vivo*, receptor agonists or antagonists may find application in the treatment of neural, heart and/or retinal diseases. A variety of suitable assays are known in the art. These assays may be based on the detection of a biological response in a target cell.

In one embodiment, binding of ligand (*e.g.*, EPO or carbamylated EPO) to the tissue protective cytokine receptor complex may be assayed in intact cells in animal models. A labeled ligand (*e.g.*, EPO or carbamylated EPO) may be administered directly to an animal, with and without a test compound. A downstream effect of the ligand (*e.g.*, EPO or carbamylated EPO) binding the tissue protective cytokine receptor complex, *e.g.*, tissue protection or cell proliferation, may be measured in the presence and the absence of test compound. For these assays, host cells to which the test compound is added may be genetically engineered to express the tissue protective cytokine receptor complex and/or ligand (*e.g.*, EPO or carbamylated EPO), which may be transient, induced or

constitutive, or stable. For the purposes of the screening methods of the present invention, a wide variety of host cells may be used including, but not limited to, tissue culture cells, mammalian cells, and yeast cells. Mammalian cells such as brain cells or other cells that express the tissue protective cytokine receptor complex may be a preferred cell type in which to carry out the assays of the present invention.

#### 5.2.2.1 Cells

In a preferred embodiment, the cell used in the methods of the invention is a mammalian cell. In a more preferred embodiment, the host cell is a human cell. In another embodiment, the host cells are primary cells isolated from a tissue or other biological sample of interest. One skilled in the art would realize that different cell types can be used in the methods of the invention for screening or identify compounds. Cell types used may endogenously express an EPO receptor, a  $\beta$  common receptor, both an EPO receptor and a  $\beta$  common receptor, or no EPO receptor or  $\beta$  common receptor. In a preferred embodiment the host cell is derived from an excitable tissue. In a preferred embodiment the host cell is derived from an excitable neuronal tissue.

By way of non-limiting examples, a responsive or excitable cells may be neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, pancreas, bone, skin, or endometrial cells or tissue. Further, non-limiting examples of responsive or excitable cells include photoreceptor (rods and cones), ganglion, bipolar, horizontal, amacrine, Müller, Purkinje, myocardium, pace maker, sinoatrial node, sinus node, junction tissue, atrioventricular node, bundle of His, hepatocytes, stellate, Kupffer, mesangial, renal epithelial, tubular interstitial, goblet, intestinal gland (crypts), enteral endocrine, glomerulosa, fasciculate, reticularis, chromaffin, pericyte, Leydig, Sertoli, sperm, Graffian follicle, primordial follicle, islets of Langerhans,  $\alpha$ -cells,  $\beta$ -cells,  $\gamma$ -cells, F-cells,

osteoprogenitor, osteoclasts, osteoblasts, endometrial stroma, endometrial, stem and endothelial cells.

Mammalian cells suitable for use in expressing the tissue protective cytokine receptor complex and transducing a receptor-mediated signal include cells that

5 express at least one of the receptor subunits used to form the functional tissue protective receptor complex. These subunits may include those of the class I cytokine receptors. In certain embodiments of the invention, it is also preferred to use a cell from the same species as the receptor to be expressed. Within a preferred embodiment, the cell is dependent upon an exogenously supplied hematopoietic growth factor for its

10 proliferation. In certain preferred embodiments, cell lines of this type are the human TF-1 cell line (ATCC number CRL-2003) and the AML-193 cell line (ATCC number CRL-9589), which are GM-CSF-dependent human leukemic cell lines. In an alternative embodiment, suitable host cells can be engineered to produce the necessary receptor subunit or other cellular component needed for the desired cellular response. For

15 example, the murine cell line BaF3 (Palacios and Steinmetz, 1985, Cell 41:727-734; Mathey-Prevot *et al.*, 1986, Mol. Cell. Biol. 6:4133-4135) or a baby hamster kidney (BHK) cell line can be transfected to express the necessary EPO-R and  $\beta_c$  receptor. The latter approach is advantageous because cell lines can be engineered to express receptor subunits from any species, thereby overcoming potential limitations arising from species

20 specificity. In the alternative, species orthologs of the human receptor cDNA can be cloned and used within cell lines from the same species, such as a mouse cDNA in the BaF3 cell line. Cell lines that are dependent upon one hematopoietic growth factor, such as GM-CSF, can thus be engineered to become dependent upon a tissue protective receptor complex ligand.

Similarly, it will be appreciated by one of ordinary skill in the art that the above assay can be performed in other cell lines that may be transformed in a similar manner. Furthermore, cell proliferation assays may be performed in cells known to harbor both EPO and  $\beta_c$  Receptor. Such cells would preferably include neuronal cell lines including, but not limited to, P-19, PC-12, and astrocytes or other cell lines known to be EPO-responsive including, but not limited to, retinal and heart cells. The presence of both EPO and  $\beta_c$  Receptors in these cell lines can be confirmed using immunoprecipitation methods well known to those of ordinary skill in the art.

Other host cells that can be used to recombinantly produce tissue protective cytokine receptor complex or used in the screening methods of the present invention include, but are not limited to, hybridomas, pre-B cells, 293 cells, 293T cells, HeLa cells, HepG2 cells, K562 cells, 3T3 cells. In a preferred embodiment, the host cells are immortalized cell lines derived from a source, *e.g.*, a tissue. Still other host cells that can be used in the present invention include, but are not limited to, yeast cells, virally-infected cells, bacteria cells, insect cells, or plant cells.

Preferred mammalian host cells include but are not limited to those derived from humans, monkeys and rodents (see, for example, Kriegler M. in "Gene Transfer and Expression: A Laboratory Manual", New York, Freeman & Co. 1990), such as monkey kidney cell line transformed by SV40 (COS-7, ATCC Accession No. CRL 1651); human embryonic kidney cell lines (293, 293-EBNA, or 293 cells subcloned for growth in suspension culture, Graham *et al.*, 1977, J. Gen. Virol., 36:59; baby hamster kidney cells (BHK, ATCC Accession No. CCL 10); chinese hamster ovary-cells-DHFR (CHO, Urlaub and Chasin, 1980, Proc. Natl. Acad. Sci. 77; 4216); mouse sertoli cells (Mather, 1980, Biol. Reprod. 23:243-251); mouse fibroblast cells (NIH-3T3), monkey kidney cells (CVI ATCC Accession No. CCL 70); african green monkey kidney cells

(VERO-76, ATCC Accession No. CRL-1587); human cervical carcinoma cells (HELA, ATCC Accession No. CCL 2); canine kidney cells (MDCK, ATCC Accession No. CCL 34); buffalo rat liver cells (BRL 3A, ATCC Accession No. CRL 1442); human lung cells (W138, ATCC Accession No. CCL 75); human liver cells (Hep G2, HB 8065); and  
5 mouse mammary tumor cells (MMT 060562, ATCC Accession No. CCL51). Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), BHK (ATCC No. CRL 1632), and BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham *et al.*, J. Gen. Virol. 36:59-72, 1977) cell lines. Additional suitable cell lines are known in the art and available from public depositories such as the American Type  
10 Culture Collection, Manassas, Va.

Other useful eukaryotic host-vector system may include yeast and insect systems. In yeast, a number of vectors containing constitutive or inducible promoters may be used with *Saccharomyces cerevisiae* (baker's yeast), *Schizosaccharomyces pombe* (fission yeast), *Pichia pastoris*, and *Hansenula polymorpha* (methylotropic  
15 yeasts). For a review see, Current Protocols in Molecular Biology, Vol. 2, 1988, Ed. Ausubel *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13; Grant *et al.*, 1987, Expression and Secretion Vectors for Yeast, in Methods in Enzymology, Eds. Wu & Grossman, 1987, Acad. Press, N.Y., Vol. 153, pp. 516-544; Glover, 1986, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, Heterologous Gene Expression  
20 in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and The Molecular Biology of the Yeast *Saccharomyces*, 1982, Eds. Strathern *et al.*, Cold Spring Harbor Press, Vols. I and II.

Fungal cells, including yeast cells, and particularly cells of the genus *Saccharomyces*, can also be used within the present invention, such as for producing  
25 receptor polypeptides, receptor fragments, polypeptide fusions, or complexes and for

cell-based assays for identifying compounds which modulate tissue protective cytokine activity. Methods for transforming yeast cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki *et al.*, U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch *et al.*, U.S. Patent No. 5,037,743; and Murray *et al.*, U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (*e.g.*, leucine). A preferred vector system for use in yeast is the POT1 vector system disclosed by Kawasaki *et al.* (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (*see, e.g.*, Kawasaki, U.S. Patent No. 4,599,311; Kingsman *et al.*, U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patent Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are also available for use in the present invention (*See, for example*, Gleeson *et al.*, 1986, J. Gen. Microbiol. 132:3459-3465, and Cregg, U.S. Patent No. 4,882,279). *Aspergillus* cells may be utilized according to the methods of McKnight *et al.*, U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino *et al.*, U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533.

Standard methods of introducing a nucleic acid sequence of interest into host cells can be used. Transformation may be by any known method for introducing

polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus and transducing a host cell with the virus, and by direct uptake of the polynucleotide. The transformation procedure used depends upon the host to be transformed. Mammalian cell transformations (*i.e.*, transfections) by direct uptake may be conducted using the calcium phosphate precipitation method of Graham & Van der Eb, 1978, *Virology* 52:546, or the various known modifications thereof. Other methods for introducing recombinant polynucleotides into cells, particularly into mammalian cells, include dextran-mediated transfection, calcium phosphate mediated transfection, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the polynucleotides into nuclei. Such methods are well-known to one of skill in the art.

Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA grown in culture, and include fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989, and Ausubel *et al.*, eds., *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., NY, 1987.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required.

Cultured mammalian cells are preferred hosts within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler *et al.*, 1978, Cell 14:725; Corsaro and Pearson, 1981, Somatic Cell Genetics 7:603; Graham and Van der Eb, 1973, Virology 52:456), electroporation (Neumann *et al.*, 1982, EMBO J. 1:841-845), DEAE-dextran mediated transfection (Ausubel *et al.*, eds., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987), and liposome-mediated transfection (Hawley-Nelson *et al.*, 1993, Focus 15:73; Ciccarone *et al.*, 1993, Focus 15:80). The production of recombinant polypeptides, including polypeptides which are components of the receptor complex of the invention, in cultured mammalian cells is disclosed, for example, by Levinson *et al.*, U.S. Patent No. 4,713,339; Hagen *et al.*, U.S. Patent No. 4,784,950; Palmiter *et al.*, U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134. In general, strong transcription promoters are preferred, such as promoters from SV-40 or cytomegalovirus. (See, *e.g.*, U.S. Patent No. 4,956,288). Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and the adenovirus major late promoter. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell.

In a preferred embodiment, stable cell lines containing the constructs of interest are generated for high throughput screening. Such stable cells lines may be generated by introducing a construct comprising a selectable marker, allowing the cells to grow for 1-2 days in an enriched medium, and then growing the cells on a selective medium. The selectable marker in the recombinant plasmid confers resistance to the



selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been inserted. Such cells are commonly referred to as "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler *et al.*, 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy *et al.*, 1980, Cell 22:817) genes can be employed in tk-, hgp<sup>r</sup>t- or ap<sup>r</sup>t- cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler *et al.*, 1980, Natl. Acad. Sci. USA 77:3567; O'Hare *et al.*, 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin *et al.*, 1981, J. Mol. Biol. 150:1); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre *et al.*, 1984, Gene 30:147) genes. A preferred selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection may be carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems may also be used to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. A preferred amplifiable selectable marker is dihydrofolate reductase, which confers

resistance to methotrexate. Other drug resistance genes (*e.g.*, multi-drug resistance, puromycin acetyltransferase) can also be used.

In certain embodiments, BaF3, an interleukin-3 dependent pre-B cell line, which possesses the  $\beta_c$  Receptor can be transfected with recombinant EPO-R. The  
5 presence of the EPO-R can then be selected by resistance to compounds such as zeocin (at 2 mg/ml) and puromycin (at 2  $\mu$ g/ml).

#### 5.2.2.2 BaF3 Cell Assays

BaF3 cell assays may also be used to identify compounds that modulate the activity of a tissue protective cytokine receptor complex.

10 For example, an assay to test for proliferation of BaF3 cells via signaling through the receptor complex can be performed as follows. In a 96 well plate eight 1:2 serial dilutions of growth medium alone (RPMI 1640, 10% fetal bovine serum, 1 mM sodium pyruvate, 2 mM L-glutamine), and the small molecule, biologic or chemical compounds of interest. The final volume of each dilution should be about 100  $\mu$ l.

15 The BaF3 parental cell line and BaF3 cells transfected with EPO-R can then be washed three times in growth media (see above), pellets resuspended in growth medium, and cells counted and diluted in growth media to 5,000 cells/100  $\mu$ l. One hundred microliters of diluted cells can then be added to each dilution of samples. The assay plate can then be incubated in a 37° C. incubator for three to four days. A 20  $\mu$ l  
20 aliquot of Alomar blue can be added to each well and the plate is incubated overnight at 37° C. The plates are read on the fluorescent plate reader at excitation wavelength of 544 and emission wavelength 590.

If the compound exhibits a tissue protective activity, the BaF3 cells should proliferate when exposed to the small molecule, biologic or chemical compound  
25 of interest.

A natural ligand for the tissue protective receptor complex can also be identified by mutagenizing a cell line expressing the receptor complex and culturing it under conditions that select for autocrine growth. (See WIPO publication WO 95/21930). Within a typical procedure, BaF3 cells expressing tissue protective receptor complex and the necessary additional subunits are mutagenized, such as with 2-ethylmethanesulfonate (EMS). The cells are then allowed to recover in the presence of IL-3, then transferred to a culture medium lacking IL-3 and IL-4. Surviving cells are screened for the production of a tissue protective receptor complex ligand, such as by adding soluble receptor to the culture medium or by assaying conditioned media on wild-type BaF3 cells and BaF3 cells expressing the receptor.

#### 5.2.2.3 Reporter Gene Assays

In certain embodiments of the invention, cells that are further engineered to express a reporter gene are used in assays to identify compounds with tissue protective activities. The reporter gene is linked to a promoter element that is responsive to the receptor-linked pathway, and the assay detects activation of transcription of the reporter gene.

For example, a reporter gene assay consistent with the methods of the invention may involve testing a compound for modulation of the activity of a tissue protective cytokine receptor complex, wherein the activated complex results in production of a transcription factor in the host cell. The host cell also comprises a SRE promoter element operably linked to a luciferase reporter gene the expression of which is regulated by the transcriptional activator, such that the luciferase gene product is either turned on or off (*i.e.* present or absent, or present in differing amounts) in the cell in response to the presence of a compound that activates the tissue protective cytokine receptor complex.

The invention yet further provides for a method for identifying a compound that modulates a tissue protective activity, comprising contacting a test compound with a tissue protective cytokine receptor complex-expressing cell, wherein said cell is transformed with a nucleic acid comprising a nucleotide sequence that

5 encodes a reporter gene operably linked to a regulatory element associated with a tissue protective cytokine receptor complex activity, identifying a test compound that increases or decreases the level of reporter gene expression relative to the level of reporter gene expression measured in the absence of the test compound, and assaying the identified test compound for a tissue protective activity. In certain embodiments of the methods of the

10 invention, the regulatory element is a serum response element. The invention still further provides for a method of identifying a compound that modulates a tissue protective activity, comprising contacting a test compound with a cell comprising, (i) a nucleic acid sequence comprising a reporter gene operably linked to a binding site specific for a DNA binding domain of a transcriptional activator, (ii) a first fusion protein comprising (A)

15 the DNA binding domain of the transcriptional activator, and (B) a first tissue protective cytokine receptor polypeptide or a fragment thereof, and (iii) a second fusion protein comprising (A) an activation domain of the transcriptional activator and (B) a second tissue protective cytokine receptor, detecting reporter gene expression, such that if the reporter gene expression detected in the presence of the test compound differs relative to

20 the reporter gene expression detected in the absence of the test compound, a compound that modulates a tissue protective activity is identified.

Any reporter gene well-known to one of skill in the art may be used in reporter gene constructs to ascertain the effect of a compound on tissue protective cytokine mediated processes or upregulation of protective globulins. A preferred

25 promoter element in this regard is a serum response element, or SRE (see, *e.g.*, Shaw *et*

*al.*, 1989, Cell 56:563-572,). A preferred form of such reporter gene is a luciferase gene (*e.g.*, firefly luciferase, renilla luciferase, and click beetle luciferase) (de Wet *et al.*, 1987, Mol. Cell. Biol. 7:725,). Expression of the luciferase gene is detected by luminescence using methods known in the art (*e.g.*, Baumgartner *et al.*, 1994, J. Biol. Chem.

5 269:29094-29101; Schenborn and Goiffin, 1993, Promega Notes 41:11). Luciferase activity assay kits are commercially available from, for example, Promega Corp., Madison, Wis. Target cell lines of this type can be used to screen libraries of chemicals, cell-conditioned culture media, fungal broths, soil samples, water samples, and the like. For example, a bank of cell-conditioned media samples can be assayed on a target cell to  
10 identify cells that produce ligand. Positive cells are then used to produce a cDNA library in a mammalian expression vector, which is divided into pools, transfected into host cells, and expressed. Media samples from the transfected cells are then assayed, with subsequent division of pools, re-transfection, subculturing, and re-assay of positive cells to isolate a cloned cDNA encoding the ligand.

15 Examples of reporter genes include, but are not limited to, fluorescent protein (*e.g.*, green fluorescent protein ("GFP"), yellow fluorescent protein, red fluorescent protein, cyan fluorescent protein, and blue fluorescent protein), beta-galactosidase ("beta-gal"), beta-glucuronidase, beta-lactamase, chloramphenicol acetyltransferase ("CAT"), secreted alkaline phosphatase ("SEAP"), horseradish  
20 peroxidase ("HRP") and alkaline phosphatase ("AP"). Alternatively, a reporter gene can also be a protein tag, such as, but not limited to, myc, His, FLAG, or GST, so that nonsense suppression will produce the peptide and the protein can be monitored by an ELISA, a western blot, or any other immunoassay to detect the protein tag. Such methods are well known to one of skill in the art.

The screening methods of the invention also encompass use of other reporter genes. Reporter genes may be obtained and the nucleotide sequence of the elements determined by any method well-known to one of skill in the art. The nucleotide sequence of a reporter gene can be obtained, *e.g.*, from the literature or a database such as GenBank. Alternatively, a polynucleotide encoding a reporter gene may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular reporter gene is not available, but the sequence of the reporter gene is known, a nucleic acid encoding the reporter gene may be chemically synthesized or obtained from a suitable source (*e.g.*, a cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the reporter gene) by PCR amplification. Once the nucleotide sequence of a reporter gene is determined, the nucleotide sequence of the reporter gene may be manipulated using methods well-known in the art for the manipulation of nucleotide sequences, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, *etc.* (see, for example, the techniques described in Sambrook *et al.*, 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel *et al.*, eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate reporter genes having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In certain embodiments, the reporter gene assays described above may be carried out in the absence of a tissue protective cytokine receptor complex ligand. In such embodiments, the test compound identified is one that interacts with the tissue protective cytokine receptor complex. In certain other embodiments, the reporter gene assays described above may be carried out in the presence of a tissue protective cytokine

receptor complex ligand, such as EPO. In such embodiments, difference in the level of transcription of the reporter gene in the presence and absence of the test compound may be detected. The level of transcript can then be compared to that in the absence of EPO. If the difference in the level of transcript detected with and without the compound is  
5 dependent on the presence of EPO, then a compound that modulates the interaction of a tissue protective cytokine receptor complex and EPO is identified. For example, if the difference in level of transcript detected increases in the presence of a compound, wherein the increase is dependent on the presence of EPO, then an agonist of the interaction between EPO and a tissue protective cytokine receptor complex is identified.

#### 10 5.2.2.4 Yeast Two-Hybrid Assays

The "two-hybrid" system can be used to detect protein-protein interactions in the yeast *Saccharomyces cerevisiae* (Fields and Song, 1989, Nature 340:245-246; U.S. Patent No. 5,283,173 by Fields and Song). Because the interactions are screened for in yeast, the inter-molecular protein interactions detected in this system  
15 occur under physiological conditions that mimic the conditions in mammalian cells (Chien *et al.*, 1991, Proc. Natl. Acad. Sci. U.S.A. 88:9578-9581). This assay utilizes the reconstitution of a transcriptional activator like GAL4 (Johnston, 1987, Microbiol. Rev. 51:458-476) through the interaction of two protein domains that have been fused to the two functional units of a transcriptional activator, the DNA-binding domain and the  
20 activation domain. This is possible due to the bipartite nature of certain transcription factors like GAL4. Being characterized as bipartite signifies that the DNA-binding and activation functions reside in separate domains and can function in trans (Keegan *et al.*, 1986, Science 231:699-704). The reconstitution of the transcriptional activator is monitored by the activation of a reporter gene such as the lacZ gene that is under the  
25 control of a promoter that contains a binding site (Upstream Activating Sequence or

UAS) for the DNA-binding domain of the transcriptional activator. This method is most commonly used either to detect an interaction between two known proteins (Fields and Song, 1989, Nature 340:245-246) or to identify interacting proteins from a population that would bind to a known protein (Durfee *et al.*, 1993, Genes Dev. 7:555-569; Gyuris  
5 *et al.*, 1993, Cell 75:791-803; Harper *et al.*, 1993, Cell 75:805-816; Vojtek *et al.*, 1993, Cell 74:205-214). Variations on the methods described herein for yeast two-hybrid assays and other methods for yeast two-hybrid assays are well known to those of skill in the art and can be used in the screening methods of the invention.

Identification of interacting proteins by the improved yeast two hybrid  
10 system is based upon the detection of expression of a reporter gene, the transcription of which is dependent upon the reconstitution of a transcriptional regulator by the interaction of two proteins, each fused to one half of the transcriptional regulator. The "bait" (*i.e.*, the tissue protective cytokine receptor complex of the present invention or derivatives or analogs thereof) and "prey" (proteins to be tested for ability to interact  
15 with the bait) proteins are expressed as fusion proteins to a DNA binding domain, and to a transcriptional regulatory domain, respectively, or vice versa.

In a specific embodiment, recombinant biological libraries expressing random peptides can be used as the source of prey nucleic acids.

In general, proteins of the bait and prey populations are provided as fusion  
20 (chimeric) proteins (preferably by recombinant expression of a chimeric coding sequence) comprising each protein contiguous to a pre-selected sequence. For one population, the pre-selected sequence is a DNA binding domain. The DNA binding domain can be any DNA binding domain, as long as it specifically recognizes a DNA sequence within a promoter. For example, the DNA binding domain is of a  
25 transcriptional activator or inhibitor. For the other population, the pre-selected sequence



is an activator or inhibitor domain of a transcriptional activator or inhibitor, respectively. The regulatory domain alone (not as a fusion to a protein sequence) and the DNA-binding domain alone (not as a fusion to a protein sequence) preferably do not detectably interact (so as to avoid false positives in the assay). The assay system further includes a reporter gene operably linked to a promoter that contains a binding site for the DNA binding domain of the transcriptional activator (or inhibitor). Accordingly, in the present method of the present invention, binding of a tissue protective cytokine receptor complex fusion protein to a prey fusion protein leads to reconstitution of a transcriptional activator (or inhibitor) which activates (or inhibits) expression of the reporter gene. The activation (or inhibition) of transcription of the reporter gene occurs intracellularly, *e.g.*, in prokaryotic or eukaryotic cells, preferably in cell culture.

The promoter that is operably linked to the reporter gene nucleotide sequence can be a native or non-native promoter of the nucleotide sequence, and the DNA binding site(s) that are recognized by the DNA binding domain portion of the fusion protein can be native to the promoter (if the promoter normally contains such binding site(s)) or non-native to the promoter. Alternatively, the transcriptional activation binding site of the desired gene(s) can be deleted and replaced with GAL4 binding sites (Bartel *et al.*, 1993, BioTechniques 14:920-924, Chasman *et al.*, 1989, Mol. Cell. Biol. 9:4746-4749).

The activation domain and DNA binding domain used in the assay can be from a wide variety of transcriptional activator proteins, as long as these transcriptional activators have separable binding and transcriptional activation domains. For example, the GAL4 protein of *S. cerevisiae* (Ma *et al.*, 1987, Cell 48:847-853), the GCN4 protein of *S. cerevisiae* (Hope & Struhl, 1986, Cell 46:885-894), the ARD1 protein of *S. cerevisiae* (Thukral *et al.*, 1989, Mol. Cell. Biol. 9:2360-2369), and the human estrogen

receptor (Kumar *et al.*, 1987, Cell 51:941-951), have separable DNA binding and activation domains. The DNA binding domain and activation domain that are employed in the fusion proteins need not be from the same transcriptional activator. In a specific embodiment, a GAL4 or LEXA DNA binding domain is employed. In another specific embodiment, a GAL4 or herpes simplex virus VP16 (Triezenberg *et al.*, 1988, Genes Dev. 2:730-742) activation domain is employed. In a specific embodiment, amino acids 1-147 of GAL4 (Ma *et al.*, 1987, Cell 48:847-853; Ptashne *et al.*, 1990, Nature 346:329-331) is the DNA binding domain, and amino acids 411-455 of VP16 (Triezenberg *et al.*, 1988, Genes Dev. 2:730-742; Cress *et al.*, 1991, Science 251:87-90) comprise the activation domain.

In a specific embodiment, plasmids encoding the different fusion protein populations can be introduced simultaneously into a single host cell (*e.g.*, a haploid yeast cell) containing one or more reporter genes, by co-transformation, to conduct the assay for protein-protein interactions. Or, preferably, the two fusion protein populations are introduced into a single cell either by mating (*e.g.*, for yeast cells) or cell fusions (*e.g.*, of mammalian cells). In a mating type assay, conjugation of haploid yeast cells of opposite mating type that have been transformed with a binding domain fusion expression construct (preferably a plasmid) and an activation (or inhibitor) domain fusion expression construct (preferably a plasmid), respectively, will deliver both constructs into the same diploid cell. The mating type of a yeast strain may be manipulated by transformation with the HO gene (Herskowitz and Jensen, 1991, Meth. Enzymol. 194:132-146).

In a preferred embodiment, a yeast interaction mating assay is employed using two different types of host cells, strain-type a and alpha of the yeast

*Saccharomyces cerevisiae*. The host cell preferably contains at least two reporter genes,

each with one or more binding sites for the DNA-binding domain (*e.g.*, of a transcriptional activator). The activator domain and DNA binding domain are each parts of chimeric proteins formed from the two respective populations of proteins. One strain of host cells, for example the  $\alpha$  strain, contains fusions of the library of nucleotide sequences with the DNA-binding domain of a transcriptional activator, such as GAL4. The hybrid proteins expressed in this set of host cells are capable of recognizing the DNA-binding site in the promoter or enhancer region in the reporter gene construct. The second set of yeast host cells, for example, the  $\alpha$  strain, contains nucleotide sequences encoding fusions of a library of DNA sequences fused to the activation domain of a transcriptional activator.

In a specific embodiment, the present invention provides a method of detecting one or more protein-protein interactions comprising (a) recombinantly expressing in a first population of yeast cells being of a first mating type a first fusion protein containing the sequence of a tissue protective cytokine receptor complex or a fragment thereof and a DNA binding domain, wherein said first population of yeast cells contains a first nucleotide sequence operably linked to a promoter driven by one or more DNA binding sites recognized by said DNA binding domain such that an interaction of said first fusion protein with a second fusion protein, said second fusion protein comprising a transcriptional activation domain, results in increased transcription of said first nucleotide sequence; (b) negatively selecting to eliminate those yeast cells in said first population in which said increased transcription of said first nucleotide sequence occurs in the absence of said second fusion protein; (c) recombinantly expressing in a second population of yeast cells of a second mating type different from said first mating type, a plurality of said second fusion proteins, each second fusion protein comprising a sequence of a fragment, derivative or analog of a protein and an activation domain of a

transcriptional activator, in which the activation domain is the same in each said second fusion protein; (d) mating said first population of yeast cells with said second population of yeast cells to form a third population of diploid yeast cells, wherein said third population of diploid yeast cells contains a second nucleotide sequence operably linked to a promoter driven by a DNA binding site recognized by said DNA binding domain such that an interaction of said first fusion protein with said second fusion protein results in increased transcription of said second nucleotide sequence, in which the first and second nucleotide sequences can be the same or different; and (e) detecting said increased transcription of said first and/or second nucleotide sequence, thereby detecting an interaction between said first fusion protein and said second fusion protein.

In one embodiment, the invention provides for a method of identifying a compound that modulates the activity of a tissue protective cytokine receptor complex, said method comprising contacting a test compound with a cell of a modified yeast strain containing (i) a nucleotide sequence encoding a reporter gene that is operably linked to a tissue protective cytokine receptor complex-responsive promoter and (ii) expresses a tissue protective cytokine receptor complex, and determining the level of activity of a tissue protective cytokine receptor complex by measuring the level of reporter gene expression, such that if the level of reporter gene activity in the presence of the compound increases or decreases relative to the level of reporter gene activity in the absence of the compound, then a compound that modulates the activity of a tissue protective cytokine receptor complex is identified.

The receptor complexes can be used in interaction trap assays (such as, for example, that described in Gyuris *et al.*, 1993, Cell 75:791-803) to identify proteins which bind to a known protein to identify inhibitors of a binding interaction. This system is a modified two-hybrid system, as described above which uses a LEU2 reporter

gene as the first nucleotide sequence and a lacZ reporter gene as the second nucleotide sequence, so that protein-protein interactions that result in the reconstitution of the transcriptional activator system can be selected for in cells lacking leucine and to expression of  $\beta$ -galactosidase. The DNA-binding domain may be the LexA DNA-binding domain, while the activator sequence may be obtained from the B42 transcriptional activation domain (Ma and Ptashne, 1987, Cell 51:113-119). The promoters of the reporter genes contain LexA binding sequences and are activated by the reconstitution of a functional transcriptional activator. Another feature of this system is that the gene encoding the DNA-binding domain fusion protein is under the influence of an inducible promoter, such as a GAL promoter, so that confirmatory tests can be performed under inducing and non-inducing conditions.

In a specific embodiment, the present invention provides a method of detecting a compound that modulates the interaction of EPO and a tissue protective cytokine receptor complex. A first population of yeast cells comprising the first fusion protein contain the sequence of a first tissue protective cytokine receptor such as an EPO receptor, and LexA DNA-binding domain. A second population of yeast cells comprise the second fusion protein comprising a B42 transcriptional activation domain fused to the sequence of a second tissue protective cytokine receptor, such as  $\beta$  common. The LEU2 reporter gene and a lacZ reporter gene are operably linked to promoters containing LexA binding sequences, such that an interaction of said first fusion protein with a second fusion protein, results in increased transcription of said reporter genes. The populations of cells comprising the fusion proteins are grown in media lacking leucine, to select. The media contains EPO and the assay is performed in the presence and absence of a test compound. Where the level of transcription differs in the presence of a

test compound, then a compound that modulates the interaction of an EPO and a tissue protective cytokine receptor complex is identified.

Still other versions of the two-hybrid approach exist, for example, a "Contingent Replication Assay" has been reported (Nallur *et al.*, 1993, Nucleic Acids Res. 21:3867-3873; Vasavada *et al.*, 1991, Proc. Natl. Acad. Sci. USA 88:10686-10690). In this case, the reconstitution of the transcription factor in mammalian cells due to the interaction of the two fusion proteins leads to the activation of transcription of the SV40 T antigen. This antigen allows the replication of the activation domain fusion plasmids. Another modification of the two-hybrid approach using mammalian cells is the "Karyoplasmic Interaction Selection Strategy" that also uses the reconstitution of a transcriptional activator (Fearon *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:7958-7962). Reporter genes used in this case have included the gene encoding the bacterial chloramphenicol acetyl transferase, the gene for cell-surface antigen CD4, and the gene encoding resistance to Hygromycin B. In both of the mammalian systems, the transcription factor that is reconstituted is a hybrid transcriptional activator in which the DNA-binding domain is from GAL4 and the activation domain is from VP16. A transcriptional activation system has been described to isolate and catalog possible protein-protein interactions within a population, and allow the comparison of such interactions between two populations (see PCT Publication WO 97/47763 published December 18, 1997).

### 5.2.3 Ligand/Tissue Protective Cytokine Receptor Complex Assays

In the cell-free and cell-based screening methods described herein, ligands of a tissue protective cytokine receptor complex can be used in the assays to identify compounds that modulate the interaction of a tissue protective cytokine receptor complex and its ligands.

In one embodiment, the invention provides for method for identifying a compound that to a tissue protective cytokine receptor complex, comprising contacting a tissue protective cytokine receptor complex with (i) a tissue protective cytokine receptor complex ligand attached to a first label and (ii) an equivalent amount of a test compound attached to a second label under conditions conducive to binding, removing unbound material from the tissue protective cytokine receptor complex, and detecting the level of the first and second labels wherein if the second label is present the compound binds the complex and if the level of the first label decreases relative to the level of the first label where the labeled ligand is contacted with a tissue protective cytokine receptor complex under conditions conducive to binding in the absence of a test compound after removal of unbound material, then a compound that binds to a tissue protective cytokine receptor complex is identified.

In yet another embodiment, the invention provides for method for identifying a compound that modulates the binding of a tissue protective cytokine receptor complex ligand to a tissue protective cytokine receptor complex, comprising contacting a tissue protective cytokine receptor complex ligand with a tissue protective cytokine receptor complex in the presence of one or more test compounds under conditions conducive to binding, and measuring the amount of tissue protective cytokine receptor complex ligand bound to the tissue protective cytokine receptor complex, such that if the amount of bound tissue protective cytokine receptor complex ligand measured in the presence of the one or more test compounds differs from the amount of bound tissue protective cytokine receptor complex ligand measured in the absence of the one or more test compounds, then a compound that modulates the binding of a tissue protective cytokine receptor complex ligand to the tissue protective cytokine receptor complex is identified.

In certain embodiments of the methods of the invention described herein where binding of a tissue protective cytokine receptor complex contacted with a tissue protective cytokine receptor complex ligand is measured, the amount of bound tissue protective cytokine receptor complex ligand is measured using a tissue protective  
5 cytokine receptor complex ligand-specific antibody.

In certain embodiments of the methods of the invention described herein above where binding of a tissue protective cytokine receptor complex contacted with a tissue protective cytokine receptor complex ligand is measured, the tissue protective cytokine receptor complex ligand is labeled and binding of the tissue protective cytokine  
10 receptor complex ligand to the tissue protective cytokine receptor complex is measured by detecting the label attached to the tissue protective cytokine receptor complex ligand.

In certain embodiments of the methods of the invention described herein above where binding of a tissue protective cytokine receptor complex contacted with a tissue protective cytokine receptor complex ligand is measured, the tissue protective  
15 cytokine receptor complex ligand is labeled and binding of the labeled ligand to the tissue protective cytokine receptor complex is measured by detecting the label attached to the tissue protective cytokine receptor complex ligand. In related embodiments, the label is fluorescent.

In one embodiment, the invention provides for a method for identifying a  
20 compound that modulates the interaction between a tissue protective cytokine receptor complex and a tissue protective cytokine receptor complex ligand, comprising contacting a tissue protective cytokine receptor complex with one or more test compounds, and measuring the tissue protective cytokine receptor complex activity, such that if the activity measured in the presence of the one or more test compounds differs from the  
25 tissue protective cytokine receptor complex activity in the absence of the one or more



test compounds, then a compound that modulates the interaction between the tissue protective cytokine receptor complex and the tissue protective cytokine receptor complex ligand is identified.

In certain embodiments of the methods of the invention described herein, the compound binds the tissue protective cytokine receptor complex. In certain embodiments of the methods of the invention described herein above, the compound binds the tissue protective cytokine receptor complex ligand.

#### 5.2.4 High Throughput Screening Assays

The cell-based and non-cell based assays described herein can be utilized in a high throughput format to screen libraries of compounds to identify those compounds that modulate a tissue protective cytokine receptor complex activity or modulate the interaction between a tissue protective cytokine receptor complex ligand and a tissue protective cytokine receptor complex. Examples of libraries of compounds that can be used in high throughput screening assays are disclosed in section 5.4 below.

High throughput screening using labeled probes can be used to identify compounds that affect the formation of specific functional sites. Functional site profiles may be compared between cells treated with a drug and untreated or control cells to identify drugs and drug candidates. Furthermore, where a specific functional site has been associated with a tissue protective activity, probes specific for such a site may be used to determine the status of the site before and after drug treatment to identify a drug that alters the status of the functional site and would, therefore, be useful in cell, tissue, or organ protective therapy. In certain embodiments, treatment with the drug restores the status of the functional site to that observed in a control sample.

The screening assays of the present invention also encompass high throughput screens and assays to identify modulators tissue protective cytokine receptor

complex activity. In accordance with this embodiment, the systems described below may be formulated into kits. To this end, cells expressing tissue protective cytokine receptor complex, or lysates thereof, can be packaged in a variety of containers, *e.g.*, vials, tubes, microtitre well plates, bottles, and the like. Other reagents can be included  
5 in separate containers and provided with the kit; *e.g.*, positive control samples, negative control samples, buffers, cell culture media, etc. The assays of the present invention may be first optimized on a small scale (*i.e.*, in test tubes), and then scaled up for high throughput assays.

### 10 5.3 ASSAYS FOR IDENTIFYING COMPOUNDS OR TESTING COMPOUNDS IDENTIFIED IN SCREENING ASSAYS

#### 5.3.1 Biological Screens or Assays

Compounds identified to have an effect on tissue protective cytokine receptor complex activity or compounds found to modulate the interaction between a tissue protective cytokine complex and a ligand may be tested for tissue protective  
15 activity, *e.g.*, protecting cells, tissues or organs. A protective activities may be further tested using *in vitro* and *in vivo* assays. *In vitro* tests that are indicative of tissue protective activity include, for example, cell proliferation assays, cell differentiation assays, or detecting the presence of proteins or nucleic acids upregulated by tissue protective cytokine receptor complex activity, *e.g.*, nucleolin, neuroglobin, and  
20 cytoglobin. Neuroglobin, for example, may be involved in facilitating the transport or the short-term storage of oxygen. Therefore, oxygen transport or storage assays may be used as an assay to identify or screen for compounds which modulate tissue protective cytokine activity.

Neuroglobin is expressed in cells and tissues of the central nervous  
25 system in response to hypoxia or ischemia and may provide protection from injury (Sun

*et al.* 2001, PNAS 98:15306-15311; Schmid *et al.*, 2003, J. Biol. Chem. 276:1932-1935). Cytoglobin may play a similar role in protection, but is expressed in a variety of tissues at varying levels (Pesce *et al.*, 2002, EMBO 3:1146-1151). In one embodiment of the invention, the levels of an upregulated protein in a cell may be measured before and after  
5 contacting the identified test compound to a cell. In certain embodiments, the presence of a upregulated protein associated with tissue protective activity in a cell, may be used to confirm the tissue protective activities of a compound.

Nucleolin may protect cells from damage. It plays numerous roles in cells including modulation of transcription processes, sequence specific RNA-binding protein,  
10 cytokinesis, nucleogenesis, signal transduction, apoptosis induced by T-cells, chromatin remodelling, or replication. It can also function as a cell surface receptor DNA/RNA helicase, DNA-dependent ATPase, protein shuttle, transcription factor component, or transcriptional repressor (Srivastava and Pollard, 1999, FASEB J., 13:1911-1922; and Ginisty *et al.*, 1999, J. Cell Sci., 112:761-772).

15 Expression of an upregulated protein may be detected by detecting mRNA levels corresponding to the protein in a cell. The mRNA can be hybridize to a probe that specifically binds a nucleic acid encoding the upregulated protein.

Hybridization may consist of, for example, Northern blot, Southern blot, array hybridization, affinity chromatography, or *in situ* hybridization.

20 In another embodiment, rapid and quantitative assay systems for screening test compounds for their ability to activate a tissue protective cytokine receptor complex can be employed. Such assays, as described for example in U.S. Patent no. 5,763,198, detect modulation of tyrosine kinase or phosphatase activities involved in signal transduction by determining the tyrosine phosphorylation state of a protein

substrate using an anti-phosphotyrosine antibody and an antibody specific for the protein substrate. These assays may be practiced in a whole cell or cell-free system.

In another embodiment, the invention provides for a method for contacting a cell that expresses a tissue protective cytokine receptor complex and lysing  
5 the cell to determine whether the tissue protective cytokine receptor complex has been phosphorylated. In related embodiments, the tissue protective cytokine receptor complex is precipitated with an anti-EPO receptor antibody and the precipitated complex is labeled with and anti-phosphotyrosine antibody for the  $\beta_c$  receptor.

Animal model systems can be used to demonstrate the tissue protective  
10 activity of a compound or to demonstrate the safety and efficacy of the compounds identified by the screening methods of the invention described above. The compounds identified in the assays can then be tested for biological activity using animal models for a type of tissue damage, disease, condition, or syndrome of interest. These include animals engineered to contain the tissue protective cytokine receptor complex coupled to  
15 a functional readout system, such as a transgenic mouse.

Animal models that can be used to test the efficacy of the cell or tissue protective activity of an identified compound include, for example, protection against the onset of Acute Experimental Allergic Encephalomyelitis (EAE) in Lewis rats, restoration or protection from diminished cognitive function in mice after receiving brain trauma  
20 (Brines *et al.*, 2000, PNAS, 97:10295-10672), and protection from induced retinal ischemia (Rosenbaum *et al.*, 1997, *Vis. Res.* 37:3443-51). Such assays are described in further detail in PCT publication no. WO02/053580, which is incorporated by reference herein in its entirety. The *in vivo* tests described therein are directed towards administration of EPO, however, a test compound that has been identified to have an  
25 effect on the activity of a tissue protective cytokine receptor complex could be

administered in place of EPO to test for tissue protective activity. Other assays for determining tissue protective activity of a compound are well known to those of skill in the art.

The assays designed to determine the tissue protective activity of a compound identified can be performed in the presence or absence of a tissue protective cytokine receptor complex ligand depending on whether the compound identified is one that interacts with the tissue protective cytokine receptor complex or modulates the interaction of a tissue protective cytokine receptor complex ligand and a tissue protective cytokine receptor complex.

In certain embodiments additional tissue protective cytokine receptor complex ligand is added to the assay to further confirm the modulation activity of a compound on the interaction of a tissue protective cytokine receptor complex ligand and a tissue protective cytokine receptor complex.

### 5.3.2 Cell Binding Assays

Alternatively, cell binding assays utilizing the cells disclosed above in section 5.2.2.1 can be utilized as well. For example, BaF3 cells could be transfected with EPO and  $\beta_c$  receptor as discussed above. Additionally, the small molecule, biologic or chemical compound of interest should be bound to a biological marker such as a fluorescent or radiolabeled marker.

In a 96 well plate eight 1:2 serial dilutions of growth medium alone (RPMI 1640, 10% fetal bovine serum, 1 mM sodium pyruvate, 2 mM L-glutamine), and the small molecule, biologic or chemical compounds of interest. The final volume of each dilution should be about 100  $\mu$ l.

The BaF3 parental cell line and BaF3 cells transfected with EPO-R can then be washed three times in growth media (see above), pellets resuspended in growth

medium, and cells counted and diluted in growth media to 5,000 cells/100  $\mu$ l. One hundred microliters of diluted cells can then be added to each dilution of samples. The assay plate can then be incubated in a 37° C. incubator for three to four days. The cells are then washed and the plate can then be read on a fluorescent plate reader or by other  
5 suitable method to detect the biomarker attached to the compound of interest.

Similarly, a competitive assay can be utilized to determine if a compound is tissue protective. In the competitive assay, a compound known to be tissue protective including, but not limited to, tissue protective cytokines such as those disclosed in U.S. Patent Application Nos. 10/188,905 and 10/185,841, can be attached to a suitable bio  
10 marker.

In a 96 well plate eight 1:2 serial dilutions of growth medium alone (RPMI 1640, 10% fetal bovine serum, 1 mM sodium pyruvate, 2 mM L-glutamine), and the remaining wells have the tissue protective compound/biomarker, the tissue protective compound/biomarker and an excess of the small molecule, biologic, or chemical  
15 compound of interest in them. The final volume of each dilution should be about 100  $\mu$ l. Once again, the BaF3 cells are seeded into the plates as disclosed above and allowed to incubate. After an appropriate amount of time, the cells are washed and the plate can then be read on a fluorescent plate reader or by other suitable method to detect the biomarker. If the readout of the plates containing tissue protective compound/biomarker  
20 and compound of interest is less than the readout of the plates containing only the tissue protective compound/biomarker then the compound of interest is tissue protective.

### 5.3.3 Cytokine and Cell Proliferation/Differentiation Activity

A receptor complex of the present invention may be helpful in identifying tissue protective compounds (small molecule, protein, chemical agents) in cytokine, cell  
25 proliferation (either inducing or inhibiting) or cell differentiation (either inducing or

inhibiting) assays. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence these assays serve as a convenient confirmation of cytokine activity. The activity of a compound can be evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, 5 DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK. These cells are cultured in the presence or absence of a test compound, and cell proliferation is detected by, for example, measuring incorporation of tritiated thymidine or by colorimetric assay based on the metabolic 10 breakdown of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (Mosman, 1983, J. Immunol. Meth. 65:55-63). The activity of a tissue protective cytokine receptor complex may, among other means, be measured by binding of a compound or ligand to the tissue protective cytokine receptor complex, cell proliferation, cell differentiation, upregulation of proteins that have a tissue protective effect, or a 15 tissue protective activity.

#### 5.3.4 Fusion Proteins

A receptor extracellular domain can be expressed as a fusion with immunoglobulin heavy chain constant regions, typically an F<sub>c</sub> fragment, which contains two constant region domains and a hinge region but lacks the variable region. Such 20 fusions are typically secreted as multimeric molecules wherein the F<sub>c</sub> portions are disulfide bonded to each other and two receptor polypeptides are arrayed in closed proximity to each other. In accordance with the present invention, the receptor polypeptides on the fusion should include at least one EPO-R extracellular domain and one  $\beta_c$  Receptor extracellular domain. Fusions of this type can be used to purify the 25 cognate ligand from solution, as an *in vitro* assay tool, to block signals *in vitro* by

specifically titrating out ligand, and as antagonists *in vivo* by administering them parenterally to bind circulating ligand and clear it from the circulation. To purify ligand, a tissue protective receptor complex chimera is added to a sample containing the ligand (*e.g.*, cell-conditioned culture media or tissue extracts) under conditions that facilitate  
5 receptor-ligand binding (typically near-physiological temperature, pH, and ionic strength). The chimera-ligand complex is then separated by the mixture using protein A, which is immobilized on a solid support (*e.g.*, insoluble resin beads). The ligand is then eluted using conventional chemical techniques, such as with a salt or pH gradient. In the alternative, the chimera itself can be bound to a solid support, with binding and elution  
10 carried out as above. Chimeras with high binding affinity are administered parenterally (*e.g.*, by intramuscular, subcutaneous or intravenous injection). Circulating molecules bind ligand and are cleared from circulation by normal physiological processes. For use in assays, the chimeras may be bound to a support via the F<sub>c</sub> region and used in an ELISA format.

15 A preferred assay system employing a ligand-binding receptor complex fusion protein or ligand-binding receptor complex fragment uses a commercially available biosensor instrument (BIAcore<sup>TM</sup>, Pharmacia Biosensor, Piscataway, N.J.), wherein the fusion protein is immobilized onto the surface of a receptor chip. Use of this instrument is disclosed by Karlsson, 1991, J. Immunol. Methods 145:229-240, and  
20 Cunningham and Wells, 1993, J. Mol. Biol. 234:554-563. A fusion protein is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within the flow cell. A test sample is passed through the cell. If ligand is present in the sample, it will bind to the immobilized fusion protein, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon  
25 resonance of the gold film. This system allows the determination of on- and off-rates,



from which binding affinity can be calculated, and assessment of stoichiometry of binding.

Ligand-binding fusion proteins may also be used within other assay systems known in the art. Such systems include Scatchard analysis for determination of binding affinity (*see*, Scatchard, 1949, Ann. NY Acad. Sci. 51:660-672) and calorimetric assays (Cunningham *et al.*, 1991, Science 253:545-548; Cunningham *et al.*, 1991, Science 254:821-825).

A receptor ligand-binding fusion protein can also be used for purification of ligand. The fusion protein is immobilized on a solid support, such as beads of agarose, cross-linked agarose, glass, cellulosic resins, silica-based resins, polystyrene, cross-linked polyacrylamide, or like materials that are stable under the conditions of use. Methods for linking polypeptides to solid supports are known in the art, and include amine chemistry, cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, and hydrazide activation. The resulting media will generally be configured in the form of a column, and fluids containing ligand are passed through the column one or more times to allow ligand to bind to the receptor polypeptide. The ligand is then eluted using changes in salt concentration or pH to disrupt ligand-receptor binding.

### 5.3.5 Other assays

One of ordinary skill in the art will recognize that the receptor complex of the present invention has utility in several well known assays including but not limited to the Ligand-Receptor Assays disclosed in Current Protocols In Immunology, vol. 4, Chapter 18, 2001, John E. Coligan Ed. John Wiley and Sons.

If the compound exhibits a tissue protective activity, one of ordinary skill in the art would recognize that it would be beneficial to verify the result using one of the

neuroprotective and tissue protective assays known to those skilled in the art, such as, but not limited to, P-19, PC-12, TF-1, and UT-7 cell assays. Additionally, various *in vivo* models such as animal models related to spinal cord injury, ischemic stroke, peripheral nerve damage, and the heart and eyes would be helpful in further

5 characterizing the tissue protective compound isolated. Suitable *in vitro* and *in vivo* assays are disclosed in U.S. Patent Application Nos. 10/188,905 and 10/185,841.

#### 5.3.5.1 Assays Using $\beta_c$ (-/-) Knock-out Animals

The invention also relates to the use of host cells and animals genetically engineered to inhibit or “knock-out” expression of the animal’s endogenous  $\beta_c$  receptor

10 and/or express the human  $\beta_c$  receptor (or mutants thereof). Such transgenic animals may be used in methods for identifying neuroprotective pathways and compounds which activate neuroprotective pathways.

Endogenous target gene expression can also be reduced by inactivating or “knocking out” the target gene or its promoter using targeted homologous recombination

15 (*e.g.*, see Smithies et al., 1985, Nature 317, 230-234; Thomas & Capecchi, 1987, Cell 51, 503-512; Thompson et al., 1989, Cell 5, 313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional target gene (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can be

20 used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express the target gene *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches are particularly suited modifications to ES (embryonic stem) cells can be used to generate animal offspring with an inactive target gene (*e.g.*, see Thomas &

25 Capecchi, 1987 and Thompson, 1989, *supra*). However this approach can be adapted for

use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors.

Once  $\beta$ c receptor (-/-) knock-out animals have been generated, the expression of the recombinant  $\beta$ c receptor gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether knock-out of the endogenous gene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include but are not limited to Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR (reverse transcriptase PCR). Samples of  $\beta$ c receptor gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the  $\beta$ c receptor transgene product.

In one embodiment of the invention,  $\beta$ c receptor (-/-) knock-out animals are used in assays to identify pathways and/or compounds that involved in a tissue protective activity. For example, normal and  $\beta$ c receptor (-/-) knock-out animals may both be administered an agent or force known to cause tissue damage, *e.g.*, ischemia. Both animals may then be administered a compound and the tissue protective activity of the compound can be determined. Methods for determining a tissue protective activity are well known to those in the art and examples of such methods are disclosed in PCT publication no. WO02/053580, which is incorporated by reference herein in its entirety. If a tissue protective activity is observed in the normal animals but not in the  $\beta$ c receptor (-/-) knock-out animals administered the same compound, then a compound with tissue protective activities dependent on the  $\beta$ c receptor is identified. Further investigation into the identified  $\beta$ c receptor dependent tissue protective pathway may consist of determining if the compound binds to the  $\beta$ c receptor or complexes thereof, or

identifying proteins or mRNA transcripts present only in the animals exhibiting the tissue protective activity. Applicants hypothesize that secondary or redundant pathways for tissue protection may be present and that  $\beta$ c receptor (-/-) knock-out animals could be used to investigate such pathways.

5

## 5.4 COMPOUNDS

The methods of the invention for screening and identifying compounds may utilize numerous types of compounds including, but not limited to, small molecules, organic molecules, non-organic molecules, peptides, polypeptides, peptide analogs including peptides comprising non-naturally occurring amino acids, *e.g.*, D-amino acids, phosphorous analogs of amino acids, such as D-amino phosphoric acids and D-amino phosphoric acids, or amino acids having non-peptide linkages, nucleic acid analogs such as phosphorothioates and PNAs.

Compounds that can be tested and identified methods described herein can include, but are not limited to, compounds obtained from any commercial source, including Aldrich (Milwaukee, WI 53233), Sigma Chemical (St. Louis, MO), Fluka Chemie AG (Buchs, Switzerland) Fluka Chemical Corp. (Ronkonkoma, NY), Eastman Chemical Company, Fine Chemicals (Kingsport, TN), Boehringer Mannheim GmbH (Mannheim, Germany), Takasago (Rockleigh, NJ), SST Corporation (Clifton, NJ), Ferro (Zachary, LA 70791), Riedel-deHaen Aktiengesellschaft (Seelze, Germany), PPG Industries Inc., Fine Chemicals (Pittsburgh, PA 15272). Further any kind of natural products may be screened using the methods of the invention, including microbial, fungal, plant or animal extracts.

Libraries of polypeptides or proteins can also be used in the assays of the invention. For example, the test compound may be a small molecule, an organic

molecule, a non-organic molecule, or a peptide. Examples of libraries of such compounds that can be screened in accordance with the methods of the invention include, but are not limited to, peptoids; random biooligomers; diversomers such as hydantoins, benzodiazepines and dipeptides; vinyllogous polypeptides; nonpeptidal  
5 peptidomimetics; oligocarbamates; peptidyl phosphonates; peptide nucleic acid libraries; antibody libraries; carbohydrate libraries; and small molecule libraries (small organic or non-organic molecule libraries). Examples of such libraries include, but are not limited to, random peptide libraries; (see, *e.g.*, Lam *et al.*, 1991, Nature 354:82-84; Houghten *et al.*, 1991, Nature 354:84-86), and combinatorial chemistry-derived molecular library  
10 made of D- and/or L-configuration amino acids, phosphopeptides (including, but not limited to, members of random or partially degenerate, directed phosphopeptide libraries; see, *e.g.*, Songyang *et al.*, 1993, Cell 72:767-778), antibodies (including, but not limited to, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')<sub>2</sub> and FAb expression library fragments, and epitope-binding  
15 fragments thereof), and small organic or inorganic molecules.

In one embodiment of the present invention, peptide libraries may be used as a source of test compounds that can be used to screen for modulators of tissue protective cytokine receptor complex interactions, such as EPO-tissue protective cytokine receptor complex. Diversity libraries, such as random or combinatorial peptide  
20 or nonpeptide libraries can be screened for molecules that specifically bind to the tissue protective cytokine receptor complex. Many libraries are known in the art that can be used, *e.g.*, chemically synthesized libraries, recombinant (*e.g.*, phage display libraries), and *in vitro* translation-based libraries.

Examples of chemically synthesized libraries are described in Fodor *et al.*, 1991, Science 251:767-773; Houghten *et al.*, 1991, Nature 354:84-86; Lam *et al.*,  
25

1991, *Nature* 354:82-84; Medynski, 1994, *Bio/Technology* 12:709-710; Gallop *et al.*,  
1994, *J. Medicinal Chemistry* 37(9):1233-1251; Ohlmeyer *et al.*, 1993, *Proc. Natl. Acad.*  
*Sci. USA* 90:10922-10926; Erb *et al.*, 1994, *Proc. Natl. Acad. Sci. USA*  
91:11422-11426; Houghten *et al.*, 1992, *Biotechniques* 13:412; Jayawickreme *et al.*,  
5 1994, *Proc. Natl. Acad. Sci. USA* 91:1614-1618; Salmon *et al.*, 1993, *Proc. Natl. Acad.*  
*Sci. USA* 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner,  
1992, *Proc. Natl. Acad. Sci. USA* 89:5381-5383.

Examples of phage display libraries are described in Scott & Smith, 1990,  
*Science* 249:386-390; Devlin *et al.*, 1990, *Science*, 249:404-406; Christian *et al.*, 1992, *J.*  
10 *Mol. Biol.* 227:711-718; Lenstra, 1992, *J. Immunol. Meth.* 152:149-157; Kay *et al.*,  
1993, *Gene* 128:59-65; and PCT Publication No. WO 94/18318 dated August 18, 1994.

By way of examples of nonpeptide libraries, a benzodiazepine library (*see*  
*e.g.*, Bunin *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91:4708-4712) can be adapted for  
use. Peptoid libraries (Simon *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:9367-9371)  
15 can also be used. Another example of a library that can be used, in which the amide  
functionalities in peptides have been permethylated to generate a chemically transformed  
combinatorial library, is described by Ostresh *et al.* (1994, *Proc. Natl. Acad. Sci. USA*  
91:11138-11142).

Screening the libraries can be accomplished by any of a variety of  
20 commonly known methods. See, *e.g.*, the following references, which disclose screening  
of peptide libraries: Parmley & Smith, 1989, *Adv. Exp. Med. Biol.* 251:215-218; Scott  
& Smith, 1990, *Science* 249:386-390; Fowlkes *et al.*, 1992; *BioTechniques* 13:422-427;  
Oldenburg *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:5393-5397; Yu *et al.*, 1994, *Cell*  
76:933-945; Staudt *et al.*, 1988, *Science* 241:577-580; Bock *et al.*, 1992, *Nature*  
25 355:564-566; Tuerk *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:6988-6992; Ellington *et*

*al.*, 1992, Nature 355:850-852; U.S. Patent No. 5,096,815, U.S. Patent No. 5,223,409, and U.S. Patent No. 5,198,346, all to Ladner *et al.*; Rebar & Pabo, 1993, Science 263:671-673; and PCT Publication No. WO 94/18318.

In a preferred embodiment, the combinatorial libraries are small organic molecule libraries, such as, but not limited to, benzodiazepines, isoprenoids, thiazolidinones, metathiazanones, pyrrolidines, morpholino compounds, and benzodiazepines. In another embodiment, the combinatorial libraries comprise peptoids; random bio-oligomers; benzodiazepines; diversomers such as hydantoins, benzodiazepines and dipeptides; vinylogous polypeptides; nonpeptidal peptidomimetics; oligocarbamates; peptidyl phosphonates; peptide nucleic acid libraries; antibody libraries; or carbohydrate libraries. Combinatorial libraries are themselves commercially available (see, *e.g.*, ComGenex, Princeton, New Jersey; Asinex, Moscow, Ru, Tripos, Inc., St. Louis, Missouri; ChemStar, Ltd, Moscow, Russia; 3D Pharmaceuticals, Exton, Pennsylvania; Martek Biosciences, Columbia, Maryland; *etc.*).

In a preferred embodiment, the library is preselected so that the compounds of the library are more amenable for cellular uptake. For example, compounds are selected based on specific parameters such as, but not limited to, size, lipophilicity, hydrophilicity, and hydrogen bonding, which enhance the likelihood of compounds getting into the cells. In another embodiment, the compounds are analyzed by three-dimensional or four-dimensional computer computation programs.

In one embodiment, the combinatorial compound library for the methods of the present invention may be synthesized. There is a great interest in synthetic methods directed toward the creation of large collections of small organic compounds, or libraries, which could be screened for pharmacological, biological or other activity. The synthetic methods applied to create vast combinatorial libraries are performed in solution

or in the solid phase, *i.e.*, on a solid support. Solid-phase synthesis makes it easier to conduct multi-step reactions and to drive reactions to completion with high yields because excess reagents can be easily added and washed away after each reaction step. Solid-phase combinatorial synthesis also tends to improve isolation, purification and screening. However, the more traditional solution phase chemistry supports a wider variety of organic reactions than solid-phase chemistry.

Combinatorial compound libraries of the present invention may be synthesized using the apparatus described in U.S. Patent No. 6,190,619 to Kilcoin *et al.*, which is hereby incorporated by reference in its entirety. U.S. Patent No. 6,190,619 discloses a synthesis apparatus capable of holding a plurality of reaction vessels for parallel synthesis of multiple discrete compounds or for combinatorial libraries of compounds.

In one embodiment, the combinatorial compound library can be synthesized in solution. The method disclosed in U.S. Patent No. 6,194,612 to Boger *et al.*, which is hereby incorporated by reference in its entirety, features compounds useful as templates for solution phase synthesis of combinatorial libraries. The template is designed to permit reaction products to be easily purified from unreacted reactants using liquid/liquid or solid/liquid extractions. The compounds produced by combinatorial synthesis using the template will preferably be small organic molecules. Some compounds in the library may mimic the effects of non-peptides or peptides. In contrast to solid phase synthesis of combinatorial compound libraries, liquid phase synthesis does not require the use of specialized protocols for monitoring the individual steps of a multistep solid phase synthesis (Egner *et al.*, 1995, J.Org. Chem. 60:2652; Anderson *et al.*, 1995, J. Org. Chem. 60:2650; Fitch *et al.*, 1994, J. Org. Chem. 59:7955; Look *et al.*, 1994, J. Org. Chem. 49:7588; Metzger *et al.*, 1993, Angew. Chem., Int. Ed. Engl.



32:894; Youngquist *et al.*, 1994, Rapid Commun. Mass Spect. 8:77; Chu *et al.*, 1995, J. Am. Chem. Soc. 117:5419; Brummel *et al.*, 1994, Science 264:399; Stevanovic *et al.*, 1993, Bioorg. Med. Chem. Lett. 3:431).

Combinatorial compound libraries useful for the methods of the present invention can be synthesized on solid supports. In one embodiment, a split synthesis method, a protocol of separating and mixing solid supports during the synthesis, is used to synthesize a library of compounds on solid supports (see *e.g.*, Lam *et al.*, 1997, Chem. Rev. 97:41-448; Ohlmeyer *et al.*, 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926 and references cited therein). Each solid support in the final library has substantially one type of compound attached to its surface. Other methods for synthesizing combinatorial libraries on solid supports, wherein one product is attached to each support, will be known to those of skill in the art (see, *e.g.*, Nefzi *et al.*, 1997, Chem. Rev. 97:449-472).

As used herein, the term "solid support" is not limited to a specific type of solid support. Rather a large number of supports are available and are known to one skilled in the art. Solid supports include silica gels, resins, derivatized plastic films, glass beads, cotton, plastic beads, polystyrene beads, alumina gels, and polysaccharides. A suitable solid support may be selected on the basis of desired end use and suitability for various synthetic protocols. For example, for peptide synthesis, a solid support can be a resin such as p-methylbenzhydrylamine (PMBHA) resin (Peptides International, Louisville, KY), polystyrenes (*e.g.*, PAM-resin obtained from Bachem Inc., Peninsula Laboratories, *etc.*), including chloromethylpolystyrene, hydroxymethylpolystyrene and aminomethylpolystyrene, poly (dimethylacrylamide)-grafted styrene co-divinyl-benzene (*e.g.*, POLYHIPE resin, obtained from Aminotech, Canada), polyamide resin (obtained from Peninsula Laboratories), polystyrene resin grafted with polyethylene glycol (*e.g.*,

TENTAGEL or ARGOGEL, Bayer, Tübingen, Germany) polydimethylacrylamide resin (obtained from Milligen/Bioscience, California), or Sepharose (Pharmacia, Sweden).

In some embodiments of the present invention, compounds can be attached to solid supports via linkers. Linkers can be integral and part of the solid support, or they may be nonintegral that are either synthesized on the solid support or attached thereto after synthesis. Linkers are useful not only for providing points of compound attachment to the solid support, but also for allowing different groups of molecules to be cleaved from the solid support under different conditions, depending on the nature of the linker. For example, linkers can be, inter alia, electrophilically cleaved, nucleophilically cleaved, photocleavable, enzymatically cleaved, cleaved by metals, cleaved under reductive conditions or cleaved under oxidative conditions. In a preferred embodiment, the compounds are cleaved from the solid support prior to high throughput screening of the compounds.

In certain embodiments, the compound of the invention is a cytokine. Examples of cytokines that can be used in the screening methods and treatment or prevention methods of the invention include, but are not limited to, IL-1 alpha, IL-1 beta, IL-1 beta, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-11, IL-13, IL-12, IL-15, IL-18, TNF-alpha, brain-derived neurotrophic factor, bone morphogenetic protein-2, beta- nerve growth factor, ciliary neurotrophic factor, epidermal growth factor, fibroblast growth factor-acidic, fibroblast growth factor-basic, granulocyte-colony stimulating factor, granulocyte-colony stimulating factor, human growth hormone, granulocyte macrophage-colony stimulating factor, hepatocyte growth factor, interferon-alpha 2a, interferon-alpha 2b, interferon-alpha 1b, interferon-beta 1a, interferon-beta 1b, interferon-gamma, insulin like growth factor-i, insulin like growth factor-II, keratinocyte growth factor, neurotrophin-3, platelet derived growth factor-bb, stem cell factor,

necrosis factor-alpha, tumor necrosis factor-beta, vascular endothelial growth factor, leptin, or prolactin. In certain embodiments, where the compound is a cytokine, the compound is recombinantly produced. In certain embodiments, where the compound is a cytokine, the compound is human.

5                   In embodiments of the methods of the invention where the compound is a small molecule, the small molecule can be specific to an EPO receptor, such as those available from Recepton (Recepton, Inc. Mountain View, CA). Other examples of small molecules include those disclosed in Goldberg et al. (2002, J. Amer. Chem. Soc. 124:544-555) and Boger and Goldberg (2001, Bioorg. and Org. Chem. 9:557-562).

#### 10   **5.4.1.1 Modified EPO and EPO Mutants**

                  In certain embodiments of the invention, the test compound used in the assay and/or the tissue protective cytokine receptor complex ligand used in assays for identifying compounds that modulate the interaction of a tissue protective cytokine receptor complex and a ligand thereof is a chemically modified EPO, a mutant EPO, or a  
15   combination thereof. For example, U.S. Patent Application No.:10/188,905, which published as 20030072737-A1 on April 17, 2003, and U.S. Patent Application No.:10/612,665 which disclose chemically modified and/or mutant EPO and which are incorporated by reference herein in their entirety. Such chemically modified and/or mutant molecules may be used in the screening assays of the invention described herein.  
20   (Also see PCT Application No. PCT/US01/49479, U.S. Patent Application Nos. 10/188,905, and 10/185,841, which are incorporated herein by reference herein in their entirety). These tissue protective cytokines protect, maintain, enhance and/or restore the function and/or viability of erythropoietin-responsive mammalian cells, tissues and organs, which include, but are not limited to, neuronal, retinal, muscle, heart, kidney  
25   cells or tissues.

Chemically modified EPO molecules that may be used in the methods of the invention include, for example, erythropoietins that have been altered by at least one modification as compared to a native erythropoietin, and preferably as compared to native human erythropoietin. The at least one modification may be a modification of at least one amino acid of the erythropoietin molecule, or a modification of at least one carbohydrate of the erythropoietin molecule. Of course, chemically modified EPO useful for the purposes herein may have a plurality of modifications compared to a native molecule, such as multiple modifications of the amino acid portion of the molecule, multiple modifications of the carbohydrate portion of the molecule, or at least one modification of the amino acid portion of the molecule and at least one modification of the carbohydrate portion of the molecule. The chemically modified EPO molecule retains its ability of protecting, maintaining, enhancing or restoring the function or viability of responsive mammalian cells, yet one or more properties of the erythropoietin molecule unrelated to the aforementioned, desirable feature may be absent as compared to the native molecule. In a preferred embodiment, the chemically modified EPO lacks erythropoietin's effects on the bone marrow, *i.e.*, increased hematocrit (erythropoiesis), vasoconstriction (high blood pressure), increased blood pressure, hyperactivation of platelets, pro-coagulant activities, and increased production of thrombocytes. More preferably, the chemically modified EPO lack erythropoiesis; most preferably the chemically modified EPO are devoid of all of erythropoietin's effects on the bone marrow.

By way of example, the chemically modified EPO of the invention may be asialoerythropoietin. In another example, the chemically modified EPO of the invention may be erythropoietin or asialoerythropoietin that has been reacted with one or more reagents that modify one or more amino groups of amino acid residues of native

erythropoietin or asialoerythropoietin. In a preferred embodiment, the chemically modified EPO is nonerythropoietic.

In one embodiment, the chemically modified EPO is an erythropoietin that has no sialic acid moieties. In a preferred embodiment, the chemically modified EPO is asialoerythropoietin, and most preferably, human asialoerythropoietin. In another embodiment, the chemically modified EPO has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 sialic acid moieties. Such partially desialylated erythropoietins are referred to herein as hyposialoerythropoietins. They may be prepared by chemical or enzymatic modification of native erythropoietin, or may be obtained by expression in a system which either does not sialylate the molecule at all or only partially sialylates the erythropoietin. The asialoerythropoietin and hyposialoerythropoietin of the invention are embraced regardless of the means by which the molecules are prepared.

In another preferred embodiment, the chemically modified EPO comprises at least one or more modified lysine residues or a modification of the N-terminal amino group of the erythropoietin molecule, such modifications as those resulting from reaction of the lysine epsilon amino group or the N-terminal amino group with an amino-group-modifying agent or agents. The modified lysine residue or modified N-terminal amino group further may be chemically reduced. In one preferred embodiment, an erythropoietin is biotinylated, carbamylated, succinylated or acetylated at one or more lysine groups or at the N-terminus. In another preferred embodiment, the lysine is reacted with an aldehyde or reducing sugar to form an imine, which optionally is then stabilized by chemical reduction such as by using sodium cyanoborohydride to form an N-alkylated lysine residue such as glucitolyl lysine, or which in the case of reducing sugars may be stabilized by Amadori or Heyns rearrangement to form an alpha-deoxy alpha-amino sugar such as alpha-deoxy-alpha-fructosyllysine. In another preferred

embodiment, the lysine or N-terminal amino group is carbamylated (carbamoylated), such as by virtue of reaction with cyanate ion, alkyl-carbamylated, aryl-carbamylated, or aryl-thiocarbamylated with an alkyl-isocyanate, aryl-isocyanate, or aryl-isothiocyanate, respectively, or it may be acylated by a reactive alkylcarboxylic or arylcarboxylic acid derivative, such as by reaction with acetic anhydride, succinic anhydride or phthalic anhydride. At least one lysine group or the N-terminal amino group may also be trinitrophenyl modified by reaction with a trinitrobenzenesulfonic acid, or preferably with one of its salts. In another embodiment, lysine residues may be modified by reaction with a glyoxal, such as reaction with glyoxal, methylglyoxal or 3-deoxyglucosone to form the corresponding alpha-carboxyalkyl derivatives.

In another embodiment, a chemically modified EPO can be generated by modifying at least one tyrosine residue of erythropoietin by using an electrophilic reagent, such as but not limited to modification by nitration or iodination, to modify an aromatic ring position.

As noted above, a tissue protective agent useful for the purposes herein may have at least one of the aforementioned modifications, but may have more than one of the above modifications. By way of example of chemically modified EPOs with one modification to the amino acid portion of the molecule and optional modification to the carbohydrate portion of the molecule, a chemically modified EPO is carbamylerythropoietin, carbamylasialoerythropoietin, carbamylhyposialoerythropoietin, acetylerthropoietin, acetylasialoerythropoietin, acetylhyposialoerythropoietin, succinylerythropoietin, succinylasialoerythropoietin, succinylhyposialoerythropoietin, biotinylerythropoietin, biotinylasialoerythropoietin, biotinylhyposialoerythropoietin, iodoerythropoietin, iodoasialoerythropoietin, iodohyposialoerythropoietin, N-epsilon-carboxymethylerythropoietin, N-epsilon-carboxymethylerythropoietin, N-epsilon-

carboxymethylhyposialoerythropoietin, and glucitolylerythropoietin, glucitolylasialoerythropoietin, glucitolylasialohypoerythropoietin. These compounds are merely exemplary of the modified erythropoietins of the invention. The foregoing trivial names are merely representative of the modifications of the native erythropoietin molecule, and as hereinbefore described, the modification of the amino group may be on one or more epsilon amino groups of lysine residues, or the N-terminal amino group, or, in the instance of nitro- or iodo-modified erythropoietins, of one or more tyrosine residues. Any combination of the foregoing is embraced herein. The present invention also embraces chemically modified EPO, comprising one or more of the aforementioned chemical modifications.

In another aspect of the invention, a method is provided for the protecting, maintaining, enhancing or restoring the function or viability of responsive mammalian cells and their associated cells, tissues and organs, by administering an effective amount of any one or more of the aforementioned chemically modified EPOs or a compound that modulates the activity of a tissue protective cytokine receptor complex such as those compounds identified by the screening methods of the invention. In one particular aspect of the method, the responsive mammalian cells and their associated cells, tissues or organs are distal to the vasculature by virtue of a tight endothelial cell barrier. In another particular aspect, the cells, tissues, organs or other bodily parts are isolated from a mammalian body, such as those intended for transplant or reattachment. By way of non-limiting examples, the responsive cell or tissue may be neuronal, retinal, muscle, heart, lung, liver, kidney, small intestine, adrenal cortex, adrenal medulla, capillary endothelial, testes, ovary, pancreas, skin, bones, or endometrial cells or tissue. These examples of responsive cells are merely illustrative. In a particular embodiment, the responsive cell or its associated cells, tissues, or organs are not excitable cells, tissues, or

organs, or do not predominantly comprise excitable cells or tissues. In another particular embodiment, the mammalian cell, tissue or organ for which an aforementioned chemically modified EPO, or a compound that modulates the activity of a tissue protective cytokine receptor complex such as those compounds identified by the screening methods of the invention, may be administered are those that have expended or will expend a period of time under at least one condition adverse to the viability of the cell, tissue or organ. Such conditions may include traumatic *in-situ* hypoxia or metabolic dysfunction, surgically-induced *in-situ* hypoxia or metabolic dysfunction, or *in-situ* toxin exposure; the latter may be associated with chemotherapy or radiation therapy. In one embodiment, the invention protects against the adverse conditions resulting from cardio-pulmonary bypass.

The invention also provides for a screening method that uses a chemically modified EPO that is i) an erythropoietin that lacks sialic acid moieties; ii) an erythropoietin that lacks N-linked or lacks O-linked carbohydrates; iii) an erythropoietin having a reduced carbohydrate content by treatment of native erythropoietin with at least one glycosidase; iv) an erythropoietin having at least one or more oxidized carbohydrates; v) an erythropoietin comprising at least one or more oxidized carbohydrates which is chemically reduced; vi) an erythropoietin comprising at least one or more modified arginine residues; vii) an erythropoietin comprising at least one or more modified lysine residues or a modification of the N-terminal amino group of the erythropoietin molecule; viii) an erythropoietin comprising at least a modified tyrosine residue; ix) an erythropoietin comprising at least a modified aspartic acid or a glutamic acid residue; x) an erythropoietin comprising at least a modified tryptophan residue; xi) an erythropoietin having at least one amino group removed; xii) an erythropoietin



comprising at least an opening of at least one of the cystine linkages in the erythropoietin molecule; or xiii) a truncated erythropoietin.

In one embodiment, the chemically modified EPO for use in the methods of the invention is asialoerythropoietin or phenylglyoxal-erythropoietin. In another  
5 embodiment, the chemically modified EPO is capable of traversing an endothelial cell barrier. The endothelial cell barrier can be selected from the group consisting of blood-brain barrier, blood-eye barrier, blood-testis barrier, blood-ovary barrier, and blood-uterus barrier.

The invention also provides for a chemically modified EPO or a  
10 compound that modulates the activity of a tissue protective cytokine receptor complex such as those compounds identified by the screening methods of the invention for use in the methods of invention that is an erythropoietin. In preferred embodiments, the compound is asialoerythropoietin. In preferred embodiments, the compound is human asialoerythropoietin. The chemically modified EPO is preferably an erythropoietin with  
15 no N-linked carbohydrates. The chemically modified EPO is preferably an erythropoietin with no O-linked carbohydrates. In one embodiment, the chemically modified EPO is an erythropoietin treated with at least one glycosidase. In another embodiment, the chemically modified EPO is periodate-oxidized erythropoietin. The periodate-oxidized erythropoietin is preferably chemically reduced with sodium  
20 cyanoborohydride. In one embodiment, the chemically modified EPO is an erythropoietin comprising a R-glyoxal moiety on the one or more arginine residues, wherein R is aryl or alkyl moiety. The erythropoietin is preferably phenylglyoxal-erythropoietin. In another embodiment, the chemically modified EPO is an erythropoietin in which at least one arginine residue is modified by reaction with a  
25 vicinal diketone selected from the group consisting of 2,3-butanedione and

cyclohexanedione. In yet another embodiment, the chemically modified EPO of the invention is an erythropoietin in which at least one arginine residue is reacted with 3-deoxyglucosone. In still another embodiment, the chemically modified EPO is an erythropoietin molecule comprising at least one biotinylated lysine or N-terminal amino group. The erythropoietin molecule can be a biotinylated erythropoietin.

The invention also provides for a chemically modified EPO that is a glucitolyl lysine erythropoietin or a fructosyl lysine erythropoietin.

In one embodiment, the chemically modified EPO of the methods of invention is an erythropoietin having at least one carbamylated lysine residue. In another embodiment, the carbamylated erythropoietin is selected from the group consisting of alpha-N-carbamoylerythropoietin; N-epsilon-carbamoylerythropoietin; alpha-N-carbamoyl, N-epsilon-carbamoylerythropoietin; alpha-N-carbamoylasialoerythropoietin; N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoyl, N-epsilon-carbamoylasialoerythropoietin; alpha-N-carbamoylhyposialoerythropoietin; N-epsilon-carbamoylhyposialoerythropoietin; and alpha-N-carbamoyl, N-epsilon-carbamoylhyposialoerythropoietin.

In one embodiment, the chemically modified EPO of the methods of invention is an erythropoietin in which at least one lysine residue is acylated. In another embodiment, a lysine residue of said erythropoietin is acetylated. In yet another embodiment, the acetylated erythropoietin is selected from the group consisting of alpha-N-acetylerythropoietin; N-epsilon-acetylerythropoietin; alpha-N-acetyl, N-epsilon-acetylerythropoietin; alpha-N-acetylasialoerythropoietin; N-epsilon-acetylasialoerythropoietin; alpha-N-acetyl, N-epsilon-acetylasialoerythropoietin; alpha-N-acetylhyposialoerythropoietin; N-epsilon-acetylhyposialoerythropoietin; and alpha-N-acetyl, N-epsilon-acetylhyposialoerythropoietin.

In one embodiment, the chemically modified EPO of the invention is an erythropoietin comprising a succinylated lysine residue. In one embodiment, the erythropoietin is selected from the group consisting of alpha-N-succinylerythropoietin; N-epsilon-succinylerythropoietin; alpha-N-succinyl, N-epsilon-succinylerythropoietin; 5 alpha-N-succinylasialoerythropoietin; N-epsilon-succinylasialoerythropoietin; alpha-N-succinyl, N-epsilon-succinylasialoerythropoietin; alpha-N-succinylhyposialoerythropoietin; N-epsilon-succinylhyposialoerythropoietin; and alpha-N-succinyl, N-epsilon-succinylhyposialoerythropoietin.

10 In one embodiment, the chemically modified EPO of the methods of invention is an erythropoietin with at least one lysine residue modified by a 2, 4, 6-trinitrobenzenesulfonic acid salt. In one aspect of the invention, the salt is 2, 4, 6-trinitrobenzenesulfonate sodium.

15 In another embodiment, the chemically modified EPO of the methods of invention is an erythropoietin in which at least one tyrosine residue is nitrated and/or iodinated.

In yet another embodiment, the chemically modified EPO of the methods of invention is an erythropoietin in which an aspartic acid and/or glutamic acid residue is reacted with a carbodiimide followed by reaction with an amine. In one aspect of the invention the amine is glycineamide.

20 Mutant EPO molecules can also be used, for example, the tissue protective cytokine receptor complex ligand may be a mutant EPO, which may be recombinantly produced, comprising one or more altered amino acid residue between position 11 to 15 of SEQ ID NO:10 [SEQ ID NO:1], position 44 to 51 of SEQ ID NO 10 [SEQ ID NO:2], position 100-108 of SEQ ID NO [SEQ ID NO:3], or position 146-151 25 of SEQ ID NO 10 [SEQ ID NO:4].

In another embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO comprising an altered amino acid residue at one or more of the following positions of SEQ ID NO: 10: 7, 20, 21, 29, 33, 38, 42, 59, 63, 67, 70, 83, 96, 126, 142, 143, 152, 153, 155, 156, or 161.

5                   In yet another embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO comprising the amino acid sequence of SEQ ID NO: 10 with one or more of the following changes (each altered sequence has been assigned a separate sequence identification number): an alanine at residue 6 of SEQ ID NO: 10 (SEQ ID NO: 15); an alanine at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 16); a serine  
10   at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 17); an isoleucine at residue 10 of SEQ ID NO: 10 (SEQ ID NO: 18); a serine at residue 11 of SEQ ID NO: 10 (SEQ ID NO: 19); an alanine at residue 12 of SEQ ID NO: 10 (SEQ ID NO: 20); an alanine at residue 13 of SEQ ID NO: 10 (SEQ ID NO: 21); an alanine residue 14 of SEQ ID NO: 10 (SEQ ID NO: 22); a glutamic acid at residue 14 of SEQ ID NO: 10 (SEQ ID NO: 23); a glutamine  
15   at residue 14 of SEQ ID NO: 10 (SEQ ID NO: 24); an alanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 25); a phenylalanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 26); an isoleucine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 27); a glutamic acid at residue 20 of SEQ ID NO: 10 (SEQ ID NO: 28); an alanine at residue 20 of SEQ ID NO: 10 (SEQ ID NO: 29); an alanine at residue 21 of SEQ ID NO: 10 (SEQ ID NO: 30); a lysine at residue 24 of SEQ ID NO: 10 (SEQ ID NO: 31); a serine at residue 29 of  
20   SEQ ID NO: 10 (SEQ ID NO: 32); a tyrosine at residue 29 of SEQ ID NO: 10 (SEQ ID NO: 33); an asparagine at residue 30 of SEQ ID NO: 10 (SEQ ID NO: 34); a threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 35); a serine at residue 33 of SEQ ID NO: 10 (SEQ ID NO: 36); a tyrosine at residue 33 of SEQ ID NO: 10 (SEQ ID NO: 37); a  
25   lysine at residue 38 of SEQ ID NO: 10 (SEQ ID NO: 38); a lysine at residue 83 of SEQ

ID NO: 10 (SEQ ID NO: 39); an asparagine at residue 42 of SEQ ID NO: 10 (SEQ ID NO: 40); an alanine at residue 42 of SEQ ID NO: 10 (SEQ ID NO: 41); an alanine at residue 43 of SEQ ID NO: 10 (SEQ ID NO: 42); an isoleucine at residue 44 of SEQ ID NO: 10 (SEQ ID NO: 43); an aspartic acid at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 44); an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 45); an alanine at residue 46 of SEQ ID NO: 10 (SEQ ID NO: 46); an alanine at residue 47 of SEQ ID NO: 10 (SEQ ID NO: 47); an isoleucine at residue 48 of SEQ ID NO: 10 (SEQ ID NO: 48); an alanine at residue 48 of SEQ ID NO: 10 (SEQ ID NO: 49); an alanine at residue 49 of SEQ ID NO: 10 (SEQ ID NO: 50); a serine at residue 49 of SEQ ID NO: 10 (SEQ ID NO: 51); a phenylalanine at residue 51 of SEQ ID NO: 10 (SEQ ID NO: 52); an asparagine at residue 51 of SEQ ID NO: 10 (SEQ ID NO: 53); an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 54); an asparagine at residue 59 of SEQ ID NO: 10 (SEQ ID NO: 55); a threonine at residue 62 of SEQ ID NO: 10 (SEQ ID NO: 56); a serine at residue 67 of SEQ ID NO: 10 (SEQ ID NO: 57); an alanine at residue 70 of SEQ ID NO: 10 (SEQ ID NO: 58); an arginine at residue 96 of SEQ ID NO: 10 (SEQ ID NO: 59); an alanine at residue 97 of SEQ ID NO: 10 (SEQ ID NO: 60); an arginine at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 61); a glutamic acid at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 62); an alanine at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 63); a threonine at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 64); an alanine at residue 101 of SEQ ID NO: 10 (SEQ ID NO: 65); an isoleucine at residue 101 of SEQ ID NO: 10 (SEQ ID NO: 66); an alanine at residue 102 of SEQ ID NO: 10 (SEQ ID NO: 67); an alanine at residue 103 of SEQ ID NO: 10 (SEQ ID NO: 68); a glutamic acid at residue 103 of SEQ ID NO: 10 (SEQ ID NO: 69); an alanine at residue 104 of SEQ ID NO: 10 (SEQ ID NO: 70); an isoleucine at residue 104 of SEQ ID NO: 10 (SEQ ID NO: 71); an alanine at residue 105 of SEQ ID NO: 10 (SEQ ID NO: 72); an alanine at residue 106 of

SEQ ID NO: 10 (SEQ ID NO: 73); an isoleucine at residue 106 of SEQ ID NO: 10 (SEQ ID NO: 74); an alanine at residue 107 of SEQ ID NO: 10 (SEQ ID NO: 75); a leucine at residue 107 of SEQ ID NO: 10 (SEQ ID NO: 76); a lysine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 77); an alanine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 78); a  
5 serine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 79); an alanine at residue 116 of SEQ ID NO: 10 (SEQ ID NO: 80); an alanine at residue 126 of SEQ ID NO: 10 (SEQ ID NO: 81); an alanine at residue 132 of SEQ ID NO: 10 (SEQ ID NO: 82); an alanine at residue 133 of SEQ ID NO: 10 (SEQ ID NO: 83); an alanine at residue 134 of SEQ ID NO: 10 (SEQ ID NO: 84); an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 85);  
10 an isoleucine at residue 142 of SEQ ID NO: 10 (SEQ ID NO: 86); an alanine at residue 143 of SEQ ID NO: 10 (SEQ ID NO: 87); an alanine at residue 146 of SEQ ID NO: 10 (SEQ ID NO: 88); a lysine at residue 147 of SEQ ID NO: 10 (SEQ ID NO: 89); an alanine at residue 147 of SEQ ID NO: 10 (SEQ ID NO: 90); a tyrosine at residue 148 of SEQ ID NO: 10 (SEQ ID NO: 91); an alanine at residue 148 of SEQ ID NO: 10 (SEQ ID NO: 92);  
15 an alanine at residue 149 of SEQ ID NO: 10 (SEQ ID NO: 93); an alanine at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 94); a glutamic acid at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 95); an alanine at residue 151 of SEQ ID NO: 10 (SEQ ID NO: 96); an alanine at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 97); a tryptophan at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 98); an alanine at residue 153 of SEQ ID NO: 10 (SEQ ID NO: 99);  
20 an alanine at residue 154 of SEQ ID NO: 10 (SEQ ID NO: 100); an alanine at residue 155 of SEQ ID NO: 10 (SEQ ID NO: 101); an alanine at residue 158 of SEQ ID NO: 10 (SEQ ID NO: 102); a serine at residue 160 of SEQ ID NO: 10 (SEQ ID NO: 103); an alanine at residue 161 of SEQ ID NO: 10 (SEQ ID NO: 104); or an alanine at residue 162 of SEQ ID NO: 10 (SEQ ID NO: 105). In one  
25 embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO that

comprises the amino acid sequence of SEQ ID NO: 10 with one or more of the amino acid residue substitutions of SEQ ID NOs: 15-105 and 119.

In yet another embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO that comprises the amino acid sequence of SEQ ID NO: 10 with a deletion of amino acid residues 44-49 of SEQ ID NO: 10.

In still another embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO that comprises the amino acid sequence of SEQ ID NO: 10 with at least one of the following changes (each altered sequence has been assigned a separate sequence identification number): i) an aspartic acid at residue 45, and a glutamic acid at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 106); ii) an asparagine at residue 30, a threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 107); iii) an aspartic acid at residue 45, a glutamic acid at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 108); iv) a glutamic acid at residue 103, and a serine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 109); v) an alanine at residue 140 and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 110); vi) an alanine at residue 140, an alanine at residue 52, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 111); vii) an alanine at residue 97, and an alanine at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 112); viii) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 113); ix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 114); x) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, and an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 115); xi) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, an alanine at residue 140, an alanine at residue 154, a lysine at residue 24, a lysine at residue 38, a lysine at residue 83, a lysine at residue 24 and an alanine at residue 15 of

SEQ ID NO: 10 (SEQ ID NO: 116); xii) a lysine at residue 24, a lysine at residue 38, and a lysine at residue 83 SEQ ID NO: 10 (SEQ ID NO: 117); or xiii) a lysine at residue 24 and an alanine at residue 15 SEQ ID NO: 10 (SEQ ID NO: 118). In one embodiment, the tissue protective cytokine receptor complex ligand is a mutant EPO that comprises the amino acid sequence of SEQ ID NO: 10 with at least one of the following amino acid residue substitutions of SEQ ID NOs: 106-118.

The tissue protective cytokine receptor complex ligand is a mutant EPO and may comprise the amino acid sequence of SEQ ID NO: 10 with at least one of the following changes, *i.e.* substitutions, (each change or combination of changes listed has been assigned a separate sequence identification number): i) an aspartic acid at residue 45, and a glutamic acid at residue 100 of SEQ ID NO: 10 (SEQ ID NO: 106); ii) an asparagine at residue 30, a threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 107); iii) an aspartic acid at residue 45, a glutamic acid at residue 150 SEQ ID NO: 10 (SEQ ID NO: 108); iv) a glutamic acid at residue 103, and a serine at residue 108 of SEQ ID NO: 10 (SEQ ID NO: 109); v) an alanine at residue 140 and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 110); vi) an alanine at residue 140, an alanine at residue 52, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 111); vii) an alanine at residue 97, and an alanine at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 112); xiii) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45 of SEQ ID NO: 10 (SEQ ID NO: 113); ix) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, and an alanine at residue 52 of SEQ ID NO: 10 (SEQ ID NO: 114); x) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, and an alanine at residue 140 of SEQ ID NO: 10 (SEQ ID NO: 115); xi) an alanine at residue 97, an alanine at residue 152, an alanine at residue 45, an alanine at residue 52, an alanine at residue 140, an alanine at residue 154, a lysine at



residue 24, a lysine at residue 38, a lysine at residue 83, a lysine at residue 24 and an alanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 116); xii) a lysine at residue 24, a lysine at residue 38, and a lysine at residue 83 SEQ ID NO: 10 (SEQ ID NO: 117); or xiii) a lysine at residue 24 and an alanine at residue 15 SEQ ID NO: 10 (SEQ ID NO: 118).

According to another aspect of the invention, the tissue protective cytokine receptor complex ligand is a mutant EPO comprising at least one of the following amino acid residue substitutions: (each change or combination of changes listed has been assigned a separate sequence identification number): a tryptophan at residue 152 of SEQ ID NO: 10 (SEQ ID NO: 98); an alanine at residue 14 and an alanine at residue 15 of SEQ ID NO: 10 (SEQ ID NO: 119); an alanine at residue 6 of SEQ ID NO: 10 (SEQ ID NO: 15); an alanine at residue 7 of SEQ ID NO: 10 (SEQ ID NO: 16); an alanine at residue 43 of SEQ ID NO: 10 (SEQ ID NO: 42); an alanine at residue 42 of SEQ ID NO: 10 (SEQ ID NO: 41); an alanine at residue 48 of SEQ ID NO: 10 (SEQ ID NO: 49); an alanine at residue 49 of SEQ ID NO: 10 (SEQ ID NO: 50); an threonine at residue 32 of SEQ ID NO: 10 (SEQ ID NO: 35); an alanine at residue 133 of SEQ ID NO: 10 (SEQ ID NO: 83); an alanine at residue 134 of SEQ ID NO: 10 (SEQ ID NO: 84); an alanine at residue 147 of SEQ ID NO: 10 (SEQ ID NO: 90); an alanine at residue 148 of SEQ ID NO: 10 (SEQ ID NO: 92); an alanine at residue 150 of SEQ ID NO: 10 (SEQ ID NO: 94); an alanine at residue 151 of SEQ ID NO: 10 (SEQ ID NO: 96); an alanine at residue 158 of SEQ ID NO: 10 (SEQ ID NO: 102); an alanine at residue 161 of SEQ ID NO: 10 (SEQ ID NO: 104); or an alanine at residue 162 of SEQ ID NO: 10 (SEQ ID NO: 105).

#### 5.4.1.2 Antibodies

In certain embodiments of the invention the compound tested in the screening assays above is an antibody. The ligand may also be an antibody specific for a tissue protective receptor cytokine receptor complex in screening assays designed to identify compounds that modulate the interaction of a known ligand with the tissue protective receptor cytokine receptor complex. Described herein are methods for the production of such antibodies. The antibodies identified may also be useful for tissue protection. The antibodies identified may also be useful for as compounds in the methods for treatment or prevention of a disease, disorder, or condition.

Such antibodies may include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Such antibodies may be utilized in conjunction with, for example, compound screening schemes, as described above, for the evaluation of the effect of test compounds on activity of a tissue protective receptor cytokine receptor complex or a tissue protective activity.

For the production of antibodies specific to a tissue protective receptor cytokine receptor complex, various host animals may be immunized by injection with an tissue protective receptor cytokine receptor complex, or a portion thereof. An antigenic portion of tissue protective receptor cytokine receptor complex can be readily predicted by algorithms known in the art.

Host animals may include, but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as

lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as a tissue protective receptor cytokine receptor complex, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with tissue protective receptor cytokine receptor complex, or portion thereof, supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, *Nature* 256, 495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor *et al.*, 1983, *Immunology Today* 4: 72; Cole *et al.*, 1983, *Proc. Natl. Acad. Sci. USA* 80, 2026-2030), and the EBV-hybridoma technique (Cole *et al.*, 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci.*, 81: 6851-6855; Neuberger *et al.*, 1984, *Nature* 312: 604-608; Takeda *et al.*, 1985, *Nature*, 314: 452-454) by splicing

the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region (see, *e.g.*, Cabilly *et al.*, U.S. Patent No. 4,816,567; and Boss *et al.*, U.S. Patent No. 4,816,397, which are incorporated herein by reference in their entirety).

In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals (see PCT International Publication No. WO 89/12690, published December 12, 1989). According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote *et al.*, 1983, Proc. Natl. Acad. Sci. U.S.A. 80:2026-2030) or by transforming human B cells with EBV virus *in vitro* (Cole *et al.*, 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, pp. 77-96). Techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl. Acad. Sci. U.S.A. 81:6851-6855; Neuberger *et al.*, 1984, Nature 312:604-608; Takeda *et al.*, 1985, Nature 314:452-454) by splicing the genes from a mouse antibody molecule specific for a tissue protective receptor cytokine receptor complex together with genes from a human antibody molecule of appropriate biological activity can also be used; such antibodies are within the scope of this invention.

Humanized antibodies are also provided (see U.S. Patent No. 5,225,539 by Winter). An immunoglobulin light or heavy chain variable region consists of a "framework" region interrupted by three hypervariable regions, referred to as complementarity determining regions (CDRs). The extent of the framework region and CDRs have been precisely defined (see, "Sequences of Proteins of Immunological Interest", Kabat, E. *et al.*, U.S. Department of Health and Human Services (1983)).

Briefly, humanized antibodies are antibody molecules from non-human species having one or more CDRs from the non-human species and a framework region from a human immunoglobulin molecule. Such CDRS-grafted antibodies have been successfully constructed against various antigens, for example, antibodies against IL-2 receptor as  
5 described in Queen *et al.*, 1989, Proc. Natl. Acad. Sci. USA 86:10029; antibodies against the cell surface receptor CAMPATH as described in Riechmann *et al.*, 1988, Nature 332:323; antibodies against hepatitis B in Co *et al.*, 1991, Proc. Natl. Acad. Sci. USA 88:2869; as well as against viral antigens of the respiratory syncytial virus in Tempest *et al.*, 1991, Bio-Technology 9:267. Humanized antibodies are most preferred for  
10 therapeutic use in humans.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242: 423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85: 5879-5883; and Ward *et al.*, 1989, Nature 334: 544-546) can be adapted to produce single chain antibodies specific for a tissue  
15 protective receptor cytokine receptor complex, or portions thereof. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the  
20 F(ab')<sub>2</sub> fragments, which can be produced by pepsin digestion of the antibody molecule and the Fab fragments, which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed (Huse *et al.*, 1989, Science, 246: 1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a tissue protective receptor cytokine receptor complex can, in turn, be utilized to generate anti-idiotypic antibodies that “mimic” the tissue protective receptor cytokine receptor complex, using techniques well known to those skilled in the art (see, *e.g.*, Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J.

5 Immunol. 147(8):2429-2438). For example antibodies which bind to the tissue protective receptor cytokine receptor complex and competitively inhibit the binding of EPO or other known ligands to the tissue protective receptor cytokine receptor complex can be used to generate anti-idiotypes that “mimic” the binding domain of the tissue protective receptor cytokine receptor complex and, therefore, bind and neutralize tissue  
10 protective receptor cytokine receptor complex ligands. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize the native ligand, which can be followed by administration of a compound identified by the methods of the invention that provides tissue protection, which may be superior in comparison to the protection provided by a native ligand. Alternatively, antibodies to the  
15 tissue protective receptor cytokine receptor complex that can act as agonists of the tissue protective receptor cytokine receptor complex activity can be generated. Such antibodies will bind to the tissue protective receptor cytokine receptor complex and activate the signal transducing activity of the receptor complex. In addition, antibodies that act as antagonist of the tissue protective receptor cytokine receptor complex activity, *i.e.*  
20 inhibit the activation of the tissue protective receptor cytokine receptor complex would be useful as a negative control in assays for identifying compounds that modulate the activity of a tissue protective receptor cytokine receptor complex. Methods for assaying for such agonists and antagonists are described in detail in the sections above.

### 5.4.2 Characterization of the Structure of Compounds

The structure of a test compound identified by the screening methods of the invention can be determined various methods by known to those of skill in the art. For example, X-ray crystallography can be used to elucidate the structure of a compound. For a review of x-ray crystallography see, *e.g.*, Blundell *et al.* 2002, Nat Rev Drug Discov 1(1):45-54.

In certain embodiments, vibrational spectroscopy (*e.g.*, but not limited to, infrared (IR) spectroscopy or Raman spectroscopy) can be used for elucidating the structure of a compound. An improved Raman spectrometer is described in US Patent No. 5,786,893 to Fink *et al.*, which is hereby incorporated by reference. Vibrational microscopy can be measured in a spatially resolved fashion to address single beads by integration of a visible microscope and spectrometer. A microscopic infrared spectrometer is described in U.S. Patent No. 5,581,085 to Reffner *et al.*, which is hereby incorporated by reference in its entirety. An instrument that simultaneously performs a microscopic infrared and microscopic Raman analysis on a sample is described in U.S. Patent No. 5,841,139 to Sostek *et al.*, which is hereby incorporated by reference in its entirety. In one embodiment of the method, compounds are synthesized on polystyrene beads doped with chemically modified styrene monomers such that each resulting bead has a characteristic pattern of absorption lines in the vibrational (IR or Raman) spectrum, by methods including but not limited to those described by Fenniri *et al.*, 2000, J. Am. Chem. Soc. 123:8151-8152. Using methods of split-pool synthesis familiar to one of skill in the art, the library of compounds is prepared so that the spectroscopic pattern of the bead identifies one of the components of the compound on the bead. Beads that have been separated according to their ability to bind a tissue protective cytokine receptor complex can be identified by their vibrational spectrum.

Mass spectrometry (*e.g.*, electrospray ionization (“ESI”) and matrix-assisted laser desorption-ionization (“MALDI”), Fourier-transform ion cyclotron resonance (“FT-ICR”) can be used both for high-throughput screening of compounds that bind to a tissue protective cytokine receptor complex and elucidating the structure of  
5 the compound.

MALDI uses a pulsed laser for desorption of the ions and a time-of-flight analyzer, and has been used for the detection of noncovalent tRNA:amino-acyl-tRNA synthetase complexes (Gruic-Sovulj *et al.*, 1997, J. Biol. Chem. 272:32084-32091). However, covalent cross-linking between the target nucleic acid and the compound is  
10 usually required for detection, since a non-covalently bound complex may dissociate during the MALDI process.

ESI mass spectrometry (“ESI-MS”) has been of greater utility for studying non-covalent molecular interactions because, unlike the MALDI process, ESI-MS generates molecular ions with little to no fragmentation (Xavier *et al.*, 2000, Trends  
15 Biotechnol. 18(8):349-356). ESI-MS has been used to study the complexes formed by HIV Tat peptide and protein with the TAR RNA (Sannes-Lowery *et al.*, 1997, Anal. Chem. 69:5130-5135).

Fourier-transform ion cyclotron resonance (“FT-ICR”) mass spectrometry provides high-resolution spectra, isotope-resolved precursor ion selection, and accurate  
20 mass assignments (Xavier *et al.*, 2000, Trends Biotechnol. 18(8):349-356). FT-ICR has been used to study the interaction of aminoglycoside antibiotics with cognate and non-cognate RNAs (Hofstadler *et al.*, 1999, Anal. Chem. 71:3436-3440; Griffey *et al.*, 1999, Proc. Natl. Acad. Sci. USA 96:10129-10133). As true for all of the mass spectrometry methods discussed herein, FT-ICR does not require labeling of the tissue protective  
25 cytokine receptor complex or a compound.



An advantage of mass spectroscopy is not only the elucidation of the structure of a compound, but also the determination of the structure of the compound bound to a tissue protective cytokine receptor complex. Such information can enable the discovery of a consensus structure of a compound that specifically binds to a tissue  
5 protective cytokine receptor complex.

NMR spectroscopy methods can also be used to characterize compounds that are associated with a tissue protective cytokine receptor complex. For example, complexed molecules can be examined by qualitatively determining changes in chemical shift, specifically from distances measured using relaxation effects, and NMR-based  
10 approaches have been used in the identification of small molecule binders of protein drug targets (Xavier *et al.*, 2000, Trends Biotechnol. 18(8):349-356). The determination of structure-activity relationships ("SAR") by NMR is the first method for NMR described in which small molecules that bind adjacent subsites are identified by two-dimensional  $^1\text{H}$ - $^{15}\text{N}$  spectra of the target protein (Shuker *et al.*, 1996, Science 274:1531-1534). The  
15 signal from the bound molecule is monitored by employing line broadening, transferred NOEs and pulsed field gradient diffusion measurements (Moore, 1999, Curr. Opin. Biotechnol. 10:54-58). A strategy for lead generation by NMR using a library of small molecules has been recently described (Fejzo *et al.*, 1999, Chem. Biol. 6:755-769).

Other examples of NMR methods that can be used for the invention  
20 include, but are not limited to, one-dimensional, two-dimensional, three dimension, four dimensional NMR methods as well as correlation spectroscopy ("COSY"), and nuclear Overhauser effect ("NOE") spectroscopy. Such methods of structure determination of compounds are well known to one of skill in the art.

Similar to mass spectroscopy, an advantage of NMR is the not only the elucidation of the structure of a compound, but also the determination of the structure of the compound bound to a tissue protective cytokine receptor complex.

## 5.5 THERAPEUTIC USE

5                   One of ordinary skill in the art would recognize that compounds identified by the present assays, and are useful as therapeutics for treatment or prevention of various diseases, disorders, and conditions. One skilled in the art would also recognize that such compounds can be used to achieve modulation of a tissue protective cytokine receptor complex. In certain embodiments of the invention, modulating the activity of a  
10   tissue protective cytokine receptor complex can be achieved by a compound that enhances or inhibits the interaction of the complex with a ligand of the complex. Both *in vitro* and *in vivo* techniques that can be used for assessing the therapeutic indications of, for example, the compounds identified by the inventive assays disclosed above are disclosed in PCT Application No. PCT/US01/49479, U.S. Patent Application Nos.  
15   10/188,905 and 10/185,841, incorporated herein by reference.

                  The aforementioned compounds identified by the methods of the invention may be useful generally for the prevention, therapeutic treatment, or prophylactic treatment of human diseases or disorders of the central nervous system or peripheral nervous system which have primarily neurological or psychiatric symptoms,  
20   ophthalmic diseases, cardiovascular diseases, cardiopulmonary diseases, respiratory diseases, kidney, urinary and reproductive diseases, bone diseases, skin diseases, gastrointestinal diseases and endocrine and metabolic abnormalities. Examples of use include, but are not limited to, protection against and repair of injury resulting from trauma and resulting inflammation to the brain (ischemic stroke, blunt trauma,  
25   subarachnoid hemorrhage), spinal cord (ischemia, blunt force trauma), peripheral nerves

(sciatic nerve injury, diabetic neuropathy, carpal tunnel syndrome), retinal (macular edema), and heart (myocardial infarct, chronic heart failure). In particular, such diseases, disorders, and conditions include hypoxic conditions, which adversely affect excitable tissues, such as excitable tissues in the central nervous system tissue, peripheral nervous system tissue, or cardiac tissue or retinal tissue such as, for example, brain, heart, or retina/eye. Therefore, the compounds identified by the methods of the invention can be used to treat or prevent damage to excitable tissue resulting from hypoxic conditions in a variety of conditions and circumstances. Non-limiting examples of such conditions and circumstances are provided in the table herein below.

In the example of the protection of neuronal tissue pathologies treatable and preventable using compounds identified by the methods of the invention, such pathologies include those which result from reduced oxygenation of neuronal tissues. Any condition which reduces the availability of oxygen to neuronal tissue, resulting in stress, damage, and finally, neuronal cell death, can be treated using compounds identified by the methods of the present invention. Generally referred to as hypoxia and/or ischemia, these conditions arise from or include, but are not limited to, stroke, vascular occlusion, prenatal or postnatal oxygen deprivation, suffocation, choking, near drowning, carbon monoxide poisoning, smoke inhalation, trauma, including surgery and radiotherapy, asphyxia, epilepsy, hypoglycemia, chronic obstructive pulmonary disease, emphysema, adult respiratory distress syndrome, hypotensive shock, septic shock, anaphylactic shock, insulin shock, sickle cell crisis, cardiac arrest, dysrhythmia, nitrogen narcosis, and neurological deficits caused by heart-lung bypass procedures.

In one embodiment, for example, the compounds identified by the methods of the present invention identified using the inventive assay could be administered alone or as part of a composition to prevent injury or tissue damage

resulting from risk of injury or tissue damage prior to, during, or subsequent to a surgical procedure or a medical procedure. For example, surgical procedures may include tumor resection or aneurysm repair, medical procedures such as chemotherapy, and natural processes such as labor or delivery. Other pathologies caused by or resulting from

5 hypoglycemia which are treatable using compounds identified by the methods of the present invention include insulin overdose, also referred to as iatrogenic hyperinsulinemia, insulinoma, growth hormone deficiency, hypocortisolism, drug overdose, and certain tumors.

Other pathologies resulting from excitable neuronal tissue damage include

10 seizure disorders, such as epilepsy, convulsions, or chronic seizure disorders. Other treatable conditions and diseases include, but are not limited to, diseases such as stroke, multiple sclerosis, hypotension, cardiac arrest, Alzheimer's disease, Parkinson's disease, cerebral palsy, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism,

15 retinal ischemia, optic nerve damage resulting from glaucoma, and neuronal loss.

The specific compounds identified by the methods of the present invention may be used to treat or prevent inflammation resulting from disease conditions or various traumas, such as physically or chemically induced inflammation. Such traumas could include angitis, chronic bronchitis, pancreatitis, osteomyelitis, rheumatoid

20 arthritis, glomerulonephritis, optic neuritis, temporal arteritis, encephalitis, meningitis, transverse myelitis, dermatomyositis, polymyositis, necrotizing fasciitis, hepatitis, and necrotizing enterocolitis.

The compounds identified by the methods of the present invention may be used to treat or prevent conditions of, and damage to, retinal tissue. Such disorders

25 include, but are not limited to retinal ischemia, macular degeneration, retinal detachment,

retinitis pigmentosa, arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, and diabetic retinopathy.

In another embodiment, the compounds identified by the methods of the present invention and principles of the invention may be used to prevent or treat injury  
5 resulting from radiation damage to excitable tissue. A further utility of the compounds identified by the methods of the present invention is in the treatment of neurotoxin poisoning, such as domoic acid shellfish poisoning, neurolathyrism, and Guam disease, amyotrophic lateral sclerosis, and Parkinson's disease.

As mentioned above, the present invention is also directed to compounds  
10 identified by the methods of the present invention for use in enhancing excitable tissue function in a mammal by peripheral administration of a tissue protective cytokine as described above. Various diseases and conditions are amenable to treatment using this method, and further, this method is useful for enhancing cognitive function in the absence of any condition or disease. These uses of the present invention are describe in  
15 further detail below and include enhancement of learning and training in both human and non-human mammals.

Conditions and diseases treatable or preventable using compounds identified by the methods of the present invention directed to the central nervous system include but are not limited to mood disorders, anxiety disorders, depression, autism,  
20 attention deficit hyperactivity disorder, and cognitive dysfunction. These conditions benefit from enhancement of neuronal function. Other disorders treatable in accordance with the teachings of the present invention include sleep disruption, for example, sleep apnea and travel-related disorders; subarachnoid and aneurismal bleeds, hypotensive shock, concussive injury, septic shock, anaphylactic shock, and sequelae of various  
25 encephalitides and meningitides, for example, connective tissue disease-related

cerebritides such as lupus. Other uses include prevention of or protection from poisoning by neurotoxins, such as domoic acid shellfish poisoning, neuroleptanesthesia, and Guam disease, amyotrophic lateral sclerosis, Parkinson's disease; postoperative treatment for embolic or ischemic injury; whole brain irradiation; sickle cell crisis; and eclampsia.

A further group of conditions treatable or preventable using compounds identified by the methods of the present invention include mitochondrial dysfunction, of either a hereditary or acquired nature, which are the cause of a variety of neurological diseases typified by neuronal injury and death. For example, Leigh disease (subacute necrotizing encephalopathy) is characterized by progressive visual loss and encephalopathy, due to neuronal drop out, and myopathy. In these cases, defective mitochondrial metabolism fails to supply enough high energy substrates to fuel the metabolism of excitable cells. An erythropoietin receptor activity modulator optimizes failing function in a variety of mitochondrial diseases. As mentioned above, hypoxic conditions adversely affect excitable tissues. The excitable tissues include, but are not limited to, central nervous system tissue, peripheral nervous system tissue, and heart tissue. In addition to the conditions described above, the compounds identified by the methods of the present invention are useful in the treatment of inhalation poisoning such as carbon monoxide and smoke inhalation, severe asthma, adult respiratory distress syndrome, and choking and near drowning. Further conditions which create hypoxic conditions or by other means induce excitable tissue damage include hypoglycemia that may occur in inappropriate dosing of insulin, or with insulin-producing neoplasms (insulinoma).

Various neuropsychologic disorders which are believed to originate from excitable tissue damage are treatable using compounds identified by the methods of the

present invention. Chronic disorders in which neuronal damage is involved and for which treatment or prevention by the present invention is provided include disorders relating to the central nervous system and/or peripheral nervous system including age-related loss of cognitive function and senile dementia, chronic seizure disorders,

5 Alzheimer's disease, Parkinson's disease, dementia, memory loss, amyotrophic lateral sclerosis, multiple sclerosis, tuberous sclerosis, Wilson's Disease, cerebral and progressive supranuclear palsy, Guam disease, Lewy body dementia, prion diseases, such as spongiform encephalopathies, *e.g.*, Creutzfeldt-Jakob disease, Huntington's disease, myotonic dystrophy, Freidrich's ataxia and other ataxias, as well as Gilles de la  
10 Tourette's syndrome, seizure disorders such as epilepsy and chronic seizure disorder, stroke, brain or spinal cord trauma, AIDS dementia, alcoholism, autism, retinal ischemia, glaucoma, autonomic function disorders such as hypertension and sleep disorders, and neuropsychiatric disorders that include, but are not limited to schizophrenia, schizoaffective disorder, attention deficit disorder, dysthymic disorder, major depressive  
15 disorder, mania, obsessive-compulsive disorder, psychoactive substance use disorders, anxiety, panic disorder, as well as unipolar and bipolar affective disorders. Additional neuropsychiatric and neurodegenerative disorders include, for example, those listed in the American Psychiatric Association's Diagnostic and Statistical manual of Mental Disorders (DSM), the most current version of which is incorporated herein by reference  
20 in its entirety.

In another embodiment, recombinant chimeric toxin molecules comprising erythropoietin can be used for therapeutic delivery of toxins to treat or prevent a proliferative disorder, such as cancer, or viral disorder, such as subacute sclerosing panencephalitis.

The following table lists additional exemplary, non-limiting indications as to the various conditions and diseases amenable to treatment by the aforementioned tissue protective cytokines.

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
Heart	Ischemia	Coronary artery disease	Acute, chronic Stable, unstable
		Myocardial infarction	Dressler's syndrome
		Angina	
		Congenital heart disease	Valvular Cardiomyopathy
		Prinzmetal angina	
		Cardiac rupture	Aneurysmatic Septal perforation
		Angiitis	
	Arrhythmia	Tachy-, bradyarrhythmia Supraventricular, ventricular Conduction abnormalities	Stable, unstable Hypersensitive carotid sinus node
	Congestive heart failure	Left, right, bi-ventricular, systolic, diastolic	Cardiomyopathies, such as idiopathic familial, infective, metabolic, storage disease, deficiencies, connective tissue disorder, infiltration and granulomas, neurovascular
		Myocarditis	Autoimmune, infective, idiopathic
		Cor pulmonale	
	Blunt and penetrating trauma		
	Toxins	Cocaine toxicity	
Vascular	Hypertension	Primary, secondary	
	Decompression sickness		
	Fibromuscular hyperplasia		
	Aneurysm	Dissecting, ruptured, enlarging	
Lungs	Obstructive	Asthma Chronic bronchitis, Emphysema and airway obstruction	
	Ischemic lung disease	Pulmonary embolism, Pulmonary thrombosis, Fat embolism	
	Environmental lung diseases		
	Ischemic lung disease	Pulmonary embolism Pulmonary thrombosis	
	Interstitial lung disease	Idiopathic pulmonary fibrosis	
	Congenital	Cystic fibrosis	
	Cor pulmonale		
	Trauma		



<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
	Pneumonia and pneumonitides	Infectious, parasitic, toxic, traumatic, burn, aspiration	
	Sarcoidosis		
Pancreas	Endocrine	Diabetes mellitus, type I and II	Beta cell failure, dysfunction Diabetic neuropathy
		Other endocrine cell failure of the pancreas	
	Exocrine	Exocrine pancreas failure	pancreatitis
Bone	Osteopenia	Primary secondary	Hypogonadism immobilisation Postmenopausal Age-related Hyperparathyroidism Hyperthyroidism Calcium, magnesium, phosphorus and/or vitamin D deficiency
	Osteomyelitis		
Skin	Avascular necrosis		
	Trauma		
	Paget's disease		
	Alopecia	Areata Totalis	Primary Secondary Male pattern baldness
		Localized generalized	Primary secondary
Autoimmune disorders	Diabetic ulceration		
	Peripheral vascular disease		
	Burn injuries		
Eye	Lupus erythematoses, Sjogren, Rheumatoid arthritis, Glomerulonephritis, Angiitis		
	Langerhan's histiocytosis		
Embryonic and fetal disorders	Optic neuritis		
	Blunt and penetrating injuries, Infections, Sarcoid, Sickle C disease, Retinal detachment, Temporal arteritis		
	Retinal ischemia, Macular degeneration, Retinitis pigmentosa, Arteriosclerotic retinopathy, Hypertensive retinopathy, Retinal artery blockage, Retinal vein blockage, Hypotension, Diabetic retinopathy, and Macular edema		
	Asphyxia		
	Ischemia		

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
CNS	Chronic fatigue syndrome, acute and chronic hypoosmolar and hyperosmolar syndromes, AIDS Dementia, Electrocutation		
	Encephalitis	Rabies, Herpes	
	Meningitis		
	Subdural hematoma		
	Nicotine addiction		
	Drug abuse and withdrawal	Cocaine, heroin, crack, marijuana, LSD, PCP, poly-drug abuse, ecstasy, opioids, sedative hypnotics, amphetamines, caffeine	
	Obsessive-compulsive disorders		
	Spinal stenosis, Transverse myelitis, Guillian Barre, Trauma, Nerve root compression, Tumoral compression, Heat stroke		
ENT	Tinnitus Meunier's syndrome Hearing loss		
	Traumatic injury, barotraumas		
Kidney	Renal failure	Acute, chronic	Vascular/ischemic, interstitial disease, diabetic kidney disease, nephrotic syndromes, infections, injury, contrast-induced, chemotherapy-induced, CPB-induced, or preventive
	Henoch S. Purpura		
Striated muscle	Autoimmune disorders	Myasthenia gravis Dermatomyositis Polymyositis	
	Myopathies	Inherited metabolic, endocrine and toxic	
	Heat stroke		
	Crush injury		
	Rhabdomyolysis		
	Mitochondrial disease		
	Infection	Necrotizing fasciitis	
Sexual dysfunction	Central and peripheral (e.g. erectile dysfunction)	Impotence secondary to medication, (diabetes)	
Liver	Hepatitis	Viral, bacterial, parasitic	
	Ischemic disease		
	Cirrhosis, fatty liver		
	Infiltrative/metabolic diseases		
Gastrointestinal	Ischemic bowel disease		

<i>Cell, tissue or organ</i>	<i>Dysfunction or pathology</i>	<i>Condition or disease</i>	<i>Type</i>
	Inflammatory bowel disease		
	Necrotizing enterocolitis		
Organ transplantation	Treatment of donor and recipient		
Reproductive tract	Infertility	Vascular Autoimmune Uterine abnormalities Implantation disorders	
Endocrine	Glandular hyper- and hypofunction		

As mentioned above, these diseases, disorders or conditions are merely illustrative of the range of benefits provided by the compounds identified by the methods of the present invention. Accordingly, this invention generally provides preventative, therapeutic, or prophylactic treatment of the consequences of mechanical trauma or of human diseases. Prevention or therapeutic or prophylactic treatment for diseases, disorders or conditions of the CNS and/or peripheral nervous system are contemplated. Prevention or therapeutic or prophylactic treatment for diseases, disorders or conditions which have a psychiatric component is provided. Prevention or therapeutic or prophylactic treatment for diseases, disorders or conditions including but not limited to those having an ophthalmic, cardiovascular, cardiopulmonary, respiratory, kidney, urinary, reproductive, gastrointestinal, endocrine, or metabolic component is provided.

In one embodiment, such a pharmaceutical composition comprising a compound that modulates the activity of a tissue protective cytokine receptor complex can be administered systemically to protect or enhance the target cells, tissue or organ. Such administration may be parenterally, via inhalation, or transmucosally, *e.g.*, orally, nasally, rectally, intravaginally, sublingually, submucosally or transdermally. Preferably, administration is parenteral, *e.g.*, via intravenous or intraperitoneal injection, and also including, but is not limited to, intra-arterial, intramuscular, intradermal and subcutaneous administration.

Selection of the preferred effective dose or amount of a compound will be readily determinable by a skilled artisan based upon considering several factors, which will be known to one of ordinary skill in the art. Such factors include the particular form of erythropoietin and/or compounds identified by the methods of the present invention, and the pharmacokinetic parameters such as bioavailability, metabolism, half-life, *etc.*, which will have been established during the usual development procedures typically employed in obtaining regulatory approval for a pharmaceutical compound. Further factors in considering the dose include the condition or disease to be treated or the benefit to be achieved in a normal individual, the body mass of the patient, the route of administration, whether administration is acute or chronic, concomitant medications, and other factors well known to affect the efficacy of administered pharmaceutical agents. Thus the precise dosage should be decided according to the judgment of the practitioner and each patient's circumstances, *e.g.*, depending upon the condition and the immune status of the individual patient, and according to standard clinical techniques.

The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention. The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

## 6. EXAMPLES

### 6.1 EXAMPLE 1: COMPETITIVE EPO BIOASSAY

The competitive binding assay described below was performed to determine whether carbamylated EPO binds to the EPO-R. UT-7 cells which express EPO-R were utilized to determine whether carbamylated EPO competitively inhibits

binding of rhEPO to the EPO-R. With binding of rhEPO to the EPO-R, cells exhibited proliferation. Inhibition of the proliferation is indicative of competitive binding.

#### Methods

For the purpose of the competitive binding assay, erythroleukemic UT-7  
5 cells (Komatsu, N. *et al.*, 1991, Cancer Res., 51:341-8) were obtained from DSMZ  
(Braunschweig, Germany, ACC137). The assays of the carbamylated EPO and rhEPO  
were performed as described in Leveque *et al.* (1996, Hematol Oncol 14:137-46) over 48  
hours. A WST-1 reduction (Roche #1 644 807) was used to quantitate the living cells.  
Signal to noise ratio of the assay was 8-15, and the half-maximal effective concentration  
10 of carbamylated EPO and rhEPO determined by a four parameter fit from concentration  
response curves using at least 6 drug concentrations. In this particular assay, the assays  
were performed with rhEPO, then separately with carbamylated EPO, and then the assay  
were performed with both rhEPO and carbamylated EPO.

The results show that cell proliferation was 0.6-fold and 1.7-fold greater,  
15 respectively, when rhEPO was administered as well as when carbamylated EPO and  
rhEPO were both incorporated into the assay. Carbamylated EPO alone, demonstrated  
no effect on cell proliferation even when used at increased concentrations.

**Figure 1** shows the concentration of the test compound (rhEPO and  
carbamylated EPO) on the x-axis, plotted against cell proliferation on the y-axis. Figure  
20 1 clearly shows that the presence of carbamylated EPO does not impact rhEPO's ability  
to stimulate the proliferation of UT-7 cells within the assay, since the competitive assay  
with both carbamylated EPO and rhEPO did not exhibit a decrease in cell proliferation,.

These results suggest that the carbamylated EPO does not bind to the  
classical EPO-R dimer complex present on UT-7 cells.

## 6.2 EXAMPLE 2: WESTERN BLOT OF RAT BRAIN MEMBRANE

Rat brain cells have been shown to be responsive to EPO. Membrane proteins of rat brain cells were isolated to determine whether these EPO responsive cells express either the  $\beta_c$  receptor or the  $\beta_c$  receptor associated with IL-3 specific receptor component.

### Methods

Rat membrane proteins were isolated using the following protocol. Rat brain was homogenized in PBS containing protease inhibitors (25  $\mu$ g/ml Leupeptin (5mg/ml in stock), 10  $\mu$ g/ml Aprotin (1mg/ml in stock), 1 mM PMSF (100 mM in stock) 20X. The homogenized tissue was then centrifuged at 3,800 rpm for 5 minutes at 4° C. The supernatant is then collected and further centrifuged at 14,000 rpm for 30 minutes at 4° C. The pellet resulting from is centrifugation is then homogenized in PBS containing the protease inhibitor and 1% Triton X-100. This homogenate is then further centrifuged at 14,000 rpm for 30 minutes at 4° C. The resulting supernatant contains the membrane proteins.

The membrane proteins were then separated by gel electrophoresis. The membrane proteins were blocked overnight with 1X casein (Vector 10X Casein Soln., Cat#: SP-5020) at 4° C. Using a Vector Avidin/Biotin Blocking Kit (Cat #: SP-2001) avidin and biotin blocks were generated. An avidin block is generated by adding 2 drops of avidin per 10 ml 1X PBS-0.1% Tween-20/1X Casein for 20 mins, rinsing the block briefly, and then washing the block with 1X PBS-0.1% Tween -20/1X Casein for 10 mins. A biotin block was generated in a similar manner, by adding 2 drops of biotin per 10 ml 1X PBS-0.1% Tween-20/1X Casein for 20 mins, rinsing the block briefly, and then washing the block with 1X PBS-0.1% Tween -20/1X Casein for 10 mins. The primary antibody (Common  $\beta$  Unit K-17 (Santa Cruz Biotechnology, Inc., Santa Cruz,

CA), dilution 1:200) was incubated for 1 hour at room temperature. The blocks were then washed four times for 10 minutes each with 1X PBS-0.1% Tween -20/1X Casein. A secondary incubation from the VECTASTAIN ABC kit (Peroxidase Rabbit IgG, Cat#: PK-4001) at a 1:1000 dilution (1 drop in 60 ml 1X PBS-0.1% Tween -20/1X Casein) was performed for 1 hour at room temperature. The block was then washed 3 times for 10 mins each with 1X PBS-0.1% Tween -20. During the wash the AB reagent was prepared in 1X PBS-0.1% Tween -20: 2 drops of reagent A of the kit into 10 ml 1X PBS-0.1% Tween -20, mix well, 2 drops of reagent B, mix well, and incubate for 30 minutes at room temperature to allow complex formation. The membranes were then incubated for 30 minutes at room temperature, then washed 3 times for 10 minutes each with 1X PBS-0.1% Tween -20. The substrate solution from the Vector Peroxidase Substrate Kit DAB in distilled water was prepared (2 drops of Buffer Stock Solution added to 10 ml distilled water, mixed, 4 drops DAB Stock Solution, mixed, 2 drops of Hydrogen Peroxide Solution, mixed) just prior to adding to the blots. The blot with the substrate was then incubated at room temperature for 10 minutes or until the color developed. The blots were then washed with 2 changes of distilled water. This procedure was also repeated using a primary antibody for the IL-3 specific subunit (T-20, Santa Cruz Biotechnology, Inc., Santa Cruz, CA).

The results indicated that both the  $\beta_c$  receptor and  $\beta_e$  receptor specific to IL-3 were identified in rat brain cell membranes.

**Figure 2** shows photographs of polyacrylamide gels of the separated membrane proteins. The first gel shows a 130 KD protein that corresponds to the  $\beta_c$  receptor. The second gel shows a 70KD protein that corresponds to the form of the  $\beta_e$  receptor specific to IL-3 receptor protein. The results clearly demonstrate the presence of the  $\beta_c$  receptor (band at 130 kD) in an erythropoietin-responsive cell (brain). This

finding suggests that a tissue protective cytokine receptor complex may form in brain cell membranes, since both EPO receptors and  $\beta_c$  receptors are present in the membranes.

### 6.3 EXAMPLE 3: IMMUNOHISTOCHEMISTRY FOR $\beta_c$ RECEPTOR IN RAT SPINAL CORD

5                   A rat spinal cord section was stained in order to test for the presence of  $\beta_c$  receptors.

#### Methods

The spinal cord of a rat was removed and fixed in paraffin. The embedded tissue was then sectioned (6  $\mu$ m). The section of spinal cord was then stained  
10   using anti- $\beta_c$  receptor antibody (K-17, Santa Cruz Biotechnology, Inc., Santa Cruz, CA.

**Figure 3** shows a photograph the sectioned spinal cord stained for  $\beta_c$  receptor using anti  $\beta_c$  receptor antibody. The stain indicates the  $\beta_c$  receptor is present within the spinal cord.

**Figure 4** shows a photograph of a close up of the stained areas of the  
15   sectioned spinal cord demonstrating reactivity with the anti- $\beta_c$  receptor antibody.

This finding suggests the presence of tissue protective cytokine receptor complexes in spinal cord tissue, since  $\beta_c$  receptors and EPO receptors are present in these tissues.

### 6.4 EXAMPLE 4: COPRECIPITATION OF EPO-R AND $\beta_c$ RECEPTOR

20                   To determine whether EPO receptor and  $\beta_c$  receptor are associated with one another in cells, receptor antibodies were used to coprecipitate complexes from cells that express both receptors. Membrane proteins from P-19 cells were separated by eletrophoresis on polyacrylamide gels and Western blots were performed to stain for EPO receptor and  $\beta_c$  receptor.

25   Methods



The immunoprecipitation of the EPO-R and  $\beta_c$  receptor from P-19 cells, neural-like embryonal carcinoma cells, was performed in accordance with the protocol outlined in Jubinsky *et al.*, 1997, (Blood 90:1867-1873). Using the antibodies specific to the receptor as described above in section 6.3 . Immunoprecipitates were then run on a  
5 polyacrylymide gel and transferred by Western and stained using antibodies specific for EPO receptors and  $\beta_c$  receptors.

**Figure 5** depicts a Western blot of the samples coimmunoprecipitated with the EPO-R and  $\beta_c$  receptor antibodies. Lane 1 shows the results of immunoprecipitation of the EPO-R in the absence of EPO on a western blot with the  $\beta_c$   
10 receptor antibody (top) and the EPO-R antibody (bottom). A band representing the  $\beta_c$  receptor can clearly be seen on the gel (arrow pointing to appropriate band, top). The arrow on the top gel, stained with  $\beta_c$  receptor antibody, indicates the position where EPO receptor migrate if it were present in the sample. Lane 2 shows the results of immunoprecipitation of the EPO-R in the presence of EPO on a western blot with the  $\beta_c$   
15 receptor antibody (top) and the EPO-R antibody (bottom). Lane 3 shows reverse immunoprecipitation, the band (bottom) indicates the presence of EPO-R in the sample immunoprecipitated using the  $\beta_c$  receptor antibody. The first arrow on the bottom gel, stained with EPO-R antibody, points to a band at about 103 kD that was identified as nucleolin (see Example 10, below). The second arrow on the bottom gel, stained with  
20 EPO-R antibody, points to a band at about 68 kD that was identified as EPO receptor (see Example 10, below).

The results suggest that EPO-R and  $\beta_c$  receptor form a complex in cells where both receptors are expressed. Finally, complexes were formed whether or not EPO itself was present (compare Lane 1 to Lane 2), suggesting that EPO is not necessary  
25 for formation of a tissue protective cytokine receptor complex.

## 6.5 EXAMPLE 5: APOPTOSIS IN CARDIOMYOCYTES WITH AND WITHOUT $\beta_c$ RECEPTOR

Apoptosis in cardiomyocytes was induced in wild type cardiomyocytes from normal mice and in cardiomyocytes from  $\beta_c$  chain knockout (-/-) mice to determine whether  $\beta_c$  receptors played a role in the protective activity of EPO on cells in preventing or delaying apoptosis.

### Methods

Cardiomyocytes were isolated from either wild type C57BL/6-129Sv or strain matched common beta chain knockout  $\beta_c$  (-/-) mice (Robb *et al.*, 1995, P.N.A.S. U.S.A. 92:9565-9569) as described (Fiordaliso *et al.*, 2001, Diabetes 50:2363-2375). Apoptotic cell death was induced by incubating in staurosporine (Sigma, 0.1  $\mu$ M) in the presence or absence of erythropoietin (100ng/ml). Wild type and  $\beta_c$  (-/-) cardiomyocyte cells served as controls (Columns 1 and 2 of Figure 7) where apoptosis was not induced. Apoptosis was induced in wild type and  $\beta_c$  (-/-) cardiomyocyte cells (Columns 3 and 4 of Figure 7), wild type cardiomyocyte cells in the presence of EPO or carbamylated EPO (Columns 5 and 6 of Figure 7), and  $\beta_c$  (-/-) cardiomyocyte cells in the presence of EPO (Column 7 of Figure 7).

Following 16 hours incubation, cells were fixed and processed for *in situ* detection of fragmented DNA using TUNEL (Roche Diagnostics).

The results are shown in **Figure 7**, which shows the percent apoptosis (y-axis) in isolated cardiomyocytes from normal and  $\beta_c$  receptor knockout mice in the presence and absence of EPO (Columns 3-7). The percent apoptosis for  $\beta_c$  (-/-) cardiomyocyte cells where apoptosis was induced in the presence of EPO (Column 7) did not significantly differ from the percent apoptosis in wild type and  $\beta_c$  (-/-) cardiomyocyte cells in the absence of EPO (see Columns 3 and 4).

The results suggest that the tissue protective activity of EPO is dependent on the presence of a  $\beta_c$  receptor in these cells.

## 6.6 EXAMPLE 6: DOWNSTREAM KINASE ACTIVATION VIA THE EPO RECEPTOR

5 To determine whether EPO and carbamylated EPO of varying concentrations were capable of inducing downstream kinase activity in cells expressing the EPO receptor, EPO receptor expressing cells were examined for the presence of an activated kinase, phosphorylated Jak2, in the presence of EPO and carbamylated EPO.

### Methods

10 BaF/3/EPOR cells were stimulated for 10 min. with EPO or carbamylated EPO at different concentrations. Activation of Jak2 was analyzed by Western blotting of cell lysates using an antibody recognizing tyrosine-phosphorylated Jak2 (PY-Jak2). Membranes were stripped and reprobed with an antibody against Jak2 to confirm equal loading.

15 **Figure 8** shows Western blots of the cell proteins separated by polyacrylamide gel electrophoresis and stained using PY-Jak2 antibodies. The top shows that phosphorylated Jak2 was present in cells stimulated with EPO at 5 nM and 50nM concentrations.

The results suggest that carbamylated EPO cannot induce downstream  
20 activation of the classical EPO receptor homodimer in cells expressing the EPO receptor. This suggests carbamylated EPO does not bind to the classical EPO receptor homodimer.

## 6.7 EXAMPLE 7: BAF3/EPOR CELL-BINDING ASSAY

To determine whether cells expressing the EPO receptor bind EPO and carbamylated EPO, BaF3 cells were contacted with each ligand and binding affinity was  
25 measured in a competitive binding assay.

### Methods

0.1 nM  $^{125}$ I EPO and graded doses of unlabelled EPO and carbamylated EPO, respectively were incubated 30 min at RT in PBS in 106 wells. The membranes were washed on filters and counted for membrane bound  $^{125}$ I EPO in a gammacounter.

- 5 The receptor bound fraction of radioligand was plotted as bound  $^{125}$ I EPO against graded doses of unlabelled ligand.

**Figure 9** shows a graph of ligand binding affinity for BaF3 cells expressing the EPO receptor. Ligand concentration in nM (x-axis) is plotted against binding (cpm) (y-axis) for BaF3 EPO receptor expressing cells contacted with EPO  
10 receptor ligands EPO and carbamylated EPO.

The BaF-3/EPO-R cells bind EPO with an affinity of about 1 nM, whereas no carbamylated EPO binding was detectable. The results indicate that carbamylated EPO does not competitively inhibit binding of EPO to cells expressing the EPO receptor.

### 15 **6.8 EXAMPLE 8: UT-7 MEMBRANE ASSAY**

To determine whether cells expressing the classical EPO receptor dimer bind EPO and carbamylated EPO, UT-7 cell membranes were contacted with each ligand and binding affinity was measured in a competitive binding assay.

### Methods

- 20 One hundred micrograms of UT-7 membranes were incubated 30 min at room temperature in 200 ul PBS with 0.1 nM  $^{125}$ I EPO and graded doses of unlabelled EPO and carbamylated EPO, respectively. The membranes were washed on filters and counted for membrane bound  $^{125}$ I EPO in a gammacounter. The receptor-bound fraction of radioligand was plotted as bound  $^{125}$ I EPO against graded doses of unlabelled ligand.

The UT-7 membranes bind EPO with an affinity of 0.1 –0.5 nM. In contrast, no carbamylated EPO binding was detected.

**Figure 10** shows a graph of ligand binding affinity for UT-7 cell membranes having the EPO receptor. Ligand concentration in nM (x-axis) is plotted against binding (cpm) (y-axis) for UT-7 EPO receptor expressing cell membranes contacted with EPO receptor ligands EPO and carbamylated EPO.

The results indicate that carbamylated EPO does not competitively inhibit binding of EPO to cells expressing both the classical EPO receptor dimer.

## 6.9 EXAMPLE 9: HEMOTOCRIT AND HEMOGLOBULIN CONCENTRATIONS

Mice were administered EPO and modified EPO to determine whether carbamylated EPO or asialoEPO have the same erythropoietic capabilities, *i.e.*, increased hematocrit and increased hemoglobin, as EPO.

### Methods

Mice were injected intravenously 5 times per week with 50µg/kg of EPO, carbamylated EPO and asialoEPO for 4 weeks. Serum hemoglobin concentrations and the hematocrite were determined by a hemoglobinometer using blood (<50 µl) withdrawn from the retroorbital plexus under isoflurane anesthesia.

**Figure 11** shows histograms of **11A**, hematocrit levels as measured by the percent volume of hematocrit (y-axis), and **11B**, hemoglobin levels measured in concentration in mM (y-axis), in mice after administration of control (vehicle), EPO, carbamylated EPO, and asialoEPO (x-axis).

The results indicate that carbamylated EPO and asialoEPO administered to mice do not induce the increased hematocrit and hemoglobin exhibited by mice which were administered EPO. This suggests that carbamylated EPO and asialoEPO interact

differently with cells *in vivo* in comparison to EPO. When combined with the experimental results of the above binding assays, these results confirm that carbamylated EPO and asialoEPO do not bind the EPO receptor and therefore cannot impart the erythropoietic activities of EPO.

## 5     **6.10 EXAMPLE 10: DETECTING THE PRESENCE OF NUCLEOLIN**

The precipitation using Western blot and the  $\beta_c$  receptor antibody described in Example 4 above was further analyzed. In particular, the protein corresponding to the 103 KD band shown in Figure 5 five Lane 1 (top) further analyzed to determine the identity of another protein that had precipitated in the samples.

### 10     Methods

P19 cells were grown to 70% confluence in complete medium. They were treated with 10 ng/ml EPO or saline (C) for 15 min, then detached by tapping on the flask, spun for 7 min at 700 rpm and resuspended in lysis buffer (TBS with protease inhibitors, 2mM CaCl<sub>2</sub>, 1% Triton and 1%NP40) to give a final concentration of 1mg protein/ml. CaCl<sub>2</sub> was present in the lysis buffer, and freezing or vortexing were avoided, to maintain protein interactions. After removal of debris by centrifugation for 10 min at 120000 rpm, the lysate was incubated with protein A sepharose (Pharmacia, 10 microL drained gel/ml) for 1h at room temperature to remove non specific binding.

The supernatant was then incubated with protein A Sepharose (10 microL gel/ml) previously coupled to the antibody for 1h and washed three times with lysis buffer. Either an antibody against beta common chain (K17, Santa Cruz Biotechnologies) or a mixture of two antibodies against EPO-R (M20 and h194, Santa Cruz Biotechnologies) at a final dilution of 1:200 were used and the incubation was run overnight at 4°C. Protein A sepharose beads were then washed five times with low detergent lysis buffer (the same as above, except 0.5% Triton and no NP40) and bound

proteins were dissociated by the addition of 30 microL of 2x Laemmli sample buffer with 5% beta-mercaptoethanol and run on a 10% SDS-PAGE.

The 103 kD band was excised from Coomassie blue stained gel, destained for few hours in 25 mM ammonium bicarbonate/40% ethanol and washed with a sequential increasing percentage of acetonitrile. Proteins were in gel-digested overnight at 30°C with trypsin (Promega, Madison WI) at a concentration of 12 ng/ml in a 25 mM ammonium bicarbonate/10% acetonitrile solution. Peptide mass fingerprinting (PMF) was performed on a Bruker ReflexIII™ matrix-assisted laser desorption/ionization (MALDI) mass spectrometer using a-cyano-4-hydroxycinnamic acid (Bruker Daltonics Billerica, MA) as a matrix. The mass spectra were externally calibrated with a mixture of 7 standard peptides in the range between 1000 to 3000 Da. Data generated were subjected to database (NCBI) searching using as programs Mascot (<http://www.matrixscience.com>) and Profound (<http://prowl.rockefeller.edu/>) allowing up to 1 missed trypsin cleavage and a mass tolerance of  $\pm 0.2$  Da. Mass spectrometry was used to determine the identity of nucleolin.

The results suggest that the P19 cells expressing the EPO receptor and the  $\beta_c$  receptor that form a tissue protective cytokine receptor complex produce nucleolin in response to administration of EPO. **Figure 12** shows a photograph of the SDS-PAGE gel with a 103 KD nucleolin protein circled.

## 6.11 EXAMPLE 11: SPINAL CORD INJURY IN $\beta_c$ RECEPTOR KNOCK-OUT MICE

Spinal cord injury was induced in  $\beta_c$  chain knockout (-/-) mice and normal (wild type) mice to further investigate the results described above in cardiomyocytes from  $\beta_c$  knockout mice and to determine whether  $\beta_c$  receptors played a role in the protective activity of EPO following spinal injury.

## Methods

Spinal cord injury was performed in mice using a modification of the protocol of Farooque (2000, *Acta Neuropathol* (Berl) 100:13-22). The technique produced an animal model that was used to test treatment of damaged excitable cells, tissues, and organs using EPO and carbamylated EPO. Specifically, C57/bl6 (wild) or  $\beta_c$  receptors knock-out (null) mice of 8-16 weeks of age were used. Each animal was confirmed null by PCR analysis of DNA obtained from the tail. Spinal injury was inflicted in five groups of mice each containing 4-9 mice. Two groups of wild type mice and three groups of null mice. During spinal injury, animals were anesthetized with isoflurane and a normal core temperature maintained. A laminectomy was performed at T3 and a 2 mm stainless steel rod was applied to the dura with 15 grams of force for 4 minutes and then removed.

Immediately following injury, a single dose of carbamylated EPO was administered to the first group of null mice, a single dose of EPO was administered to the second group of null mice, PBS was administered as a control to the third group of null mice, a single dose of carbamylated EPO was administered to the first group of wild type mice, and a single dose of PBS was administered to the second group of wild type mice.

Animals were then assessed for 42 days using the scoring system of Basso et al. (1995, *J. Neurotrauma* 12:1-21). The Basso, Beattie, Bresnahan (BBB) scale is a 21-point scale for testing normal overground motor locomotion that is commonly used to measure of functional recovery following spinal cord injury. The BBB score is very sensitive for detecting fine motor deficits and also assesses the degree of recovery after a spinal cord injury. BBB motor scores were plotted versus number of days post injury for all groups of animals. Clinical severity was then calculated measuring the area under the curve (AUC) of the plot of BBB versus number of days post injury. Clinical severity



values among groups of mice were analyzed statically for differences and a bar graph showing the clinical severity values for each group was generated.

The results confirm those in cardiomyocyte cells and suggest that the tissue protective activity of EPO is dependent on the presence of a  $\beta_c$  receptor in animals.

5 The results are shown in **Figure 13**, which shows the BBB motor score (y-axis) versus time in days following injury (x-axis). The wild type group of mice administered carbamylated EPO exhibited increased motor scores throughout the 42 post-injury time period in comparison to all other groups. The results are shown in **Figure 14**, which shows the clinical severity measured in area under curve (AUC) (y-axis) and groups of  
10 mice (x-axis). The wild type group of mice administered carbamylated EPO (column 1) exhibited a significant increase in clinical severity values (*i.e.*, less severity) in comparison to all other groups (columns 2-5). The group of null mice administered EPO (column 5) exhibited very high mortality in comparison to all other groups. The results suggest that a  $\beta_c$  receptor is necessary to achieve the therapeutic benefit of EPO in  
15 treating injury.

The invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the  
20 invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All references cited herein are incorporated by reference herein in their  
25 entireties for all purposes.

## WHAT IS CLAIMED:

1. A method for identifying a compound that modulates a tissue protective activity, comprising:
  - 5 (a) contacting a test compound with a tissue protective cytokine receptor complex;
  - (b) measuring the level of tissue protective cytokine receptor complex activity;
  - (c) identifying a test compound which increases or decreases the level of  
10 tissue protective cytokine receptor complex activity as compared to the level of tissue protective cytokine receptor complex activity measured in the absence of the test compound; and
  - (d) assaying the identified test compound for tissue protective activity.
2. The method of claim 1 wherein the tissue protective cytokine receptor complex  
15 activity is measured by measuring the binding of the test compound to the tissue protective cytokine receptor complex.
3. The method of claim 2, wherein the test compound is labeled and binding of the labeled test compound to the tissue protective cytokine receptor complex is measured by detecting the label attached to the test compound.
- 20 4. The method of claim 2 wherein the tissue protective cytokine receptor complex activity is measured by measuring the binding of the test compound to the tissue protective cytokine receptor complex.
5. A method for identifying a compound that modulates a tissue protective activity, comprising:
  - 25 (a) contacting a test compound with a tissue protective cytokine receptor complex-expressing cell; and
  - (b) measuring the level of tissue protective cytokine receptor complex activity in the cell;

- (c) identifying a test compound which increases or decreases the tissue protective cytokine receptor complex activity as compared to the level of tissue protective cytokine receptor complex activity measured in the absence of the test compound; and
- 5 (d) assaying for the identified test compound for a tissue protective activity.
6. The method of Claim 5, wherein the tissue protective cytokine receptor complex activity is measured by a cell proliferation assay.
7. The method of Claim 5, wherein the cell is recombinantly engineered to express at least one EPO or  $\beta$  common receptors.
- 10 8. The method of Claim 7, wherein the cell endogenously expresses an EPO receptor and is transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a  $\beta$  common receptor polypeptide.
9. The method of Claim 7, wherein the cell endogenously expresses a  $\beta$  common receptor and is transformed with a nucleic acid comprising a nucleotide sequence that (i)
- 15 is operably linked to a promoter, and (ii) encodes an EPO receptor polypeptide.
10. The method of Claim 8 or 9, wherein the nucleotide sequence is derived from the same species as the cell.
11. A method for identifying a compound that modulates a tissue protective activity, comprising:
- 20 (a) contacting a test compound with a cell which is recombinantly engineered to express an EPO receptor, wherein said cell is transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a  $\beta$  common receptor polypeptide;
- (b) measuring the level of tissue protective cytokine receptor complex
- 25 activity in the cell;
- (c) identifying a test compound that increases or decreases the level of tissue protective cytokine receptor complex activity in the cell relative to the level of tissue protective cytokine receptor complex activity measured in a control cell,

wherein the control cell is of the same cell type as the cell of step (a) and is not transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a  $\beta$  common receptor polypeptide; and

5 (d) assaying the identified test compound for a tissue protective activity.

12. A method for identifying a compound that modulates a tissue protective activity, comprising:

- (a) contacting a test compound with a recombinant cell that expresses a  $\beta$  common receptor, wherein said cell is transformed with a nucleic acid  
10 comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a EPO receptor polypeptide;
- (b) measuring the level of tissue protective cytokine receptor complex activity in the cell;
- (c) identifying a test compound that increases or decreases the level of tissue  
15 protective cytokine receptor complex activity in the cell relative to the level of tissue protective cytokine receptor complex activity measured in a control cell, wherein the control cell is of the same cell type as the cell of step (a) and is not transformed with a nucleic acid comprising a nucleotide sequence that (i) is operably linked to a promoter, and (ii) encodes a  $\beta$  common receptor polypeptide;  
20 and
- (d) assaying the identified test compound for a tissue protective activity.

13. A method for identifying a compound that modulates a tissue protective activity, comprising:

- (a) contacting a test compound with a tissue protective cytokine receptor  
25 complex-expressing cell, wherein said cell is transformed with a nucleic acid comprising a nucleotide sequence that encodes a reporter gene operably linked to a regulatory element associated with a tissue protective cytokine receptor complex activity;
- (b) identifying a test compound that increases or decreases the level of  
30 reporter gene expression relative to the level of reporter gene expression measured in the absence of the test compound, and
- (c) assaying the identified test compound for a tissue protective activity.

14. The method of Claim 13, wherein the regulatory element is a serum response element.

15. A method of identifying a compound that modulates a tissue protective activity, comprising:

- 5           (a)     contacting a test compound with a cell comprising:
- (i)     a nucleic acid sequence comprising a reporter gene operably linked to a binding site specific for a DNA binding domain of a transcriptional activator;
  - 10       (ii)    a first fusion protein comprising (A) the DNA binding domain of the transcriptional activator, and (B) a first tissue protective cytokine receptor polypeptide or a fragment thereof; and
  - (iii)   a second fusion protein comprising (A) an activation domain of the transcriptional activator and (B) a second tissue protective cytokine receptor,

15           (b)     detecting reporter gene expression,  
such that if the reporter gene expression in (b) differs relative to the reporter gene expression detected in the absence of the test compound, a compound that modulates a tissue protective activity is identified.

16. The method of any one of Claims 5, 11, 12, 13, or 15, wherein the cell is a  
20     prokaryotic cell.

17. The method of any one of Claims 5, 11, 12, 13, or 15, wherein the cell is a eukaryotic cell.

18. The method of Claim 17, wherein the eukaryotic cell is a human cell.

19. The method of any one of Claims 5, 11, 12, 13, or 15, wherein the cell  
25     endogenously expresses at least one receptor of the tissue protective cytokine receptor complex.

20. The method of any one of Claims 5, 11, 12, 13, or 15, wherein the cell is a BaF3 cell.

21. A method of identifying a compound that modulates the activity of a tissue protective cytokine receptor complex, said method comprising:

- 5 (a) contacting a test compound with a cell of a modified yeast strain containing (i) a nucleotide sequence encoding a reporter gene that is operably linked to a tissue protective cytokine receptor complex-responsive promoter and (ii) expresses a tissue protective cytokine receptor complex; and
- (b) determining the level of activity of a tissue protective cytokine receptor complex by measuring the level of reporter gene expression, such that if the level of reporter gene activity in the presence of the compound increases or decreases relative to the level of reporter gene activity in the absence of the compound, then a compound that modulates the activity of a tissue protective cytokine receptor complex is identified.
- 10

22. A method for identifying a compound that binds to a tissue protective cytokine receptor complex, comprising:

- 15 (a) contacting a tissue protective cytokine receptor complex with (i) a tissue protective cytokine receptor complex ligand attached to a first label and (ii) an equivalent amount of a test compound attached to a second label under conditions conducive to binding;
- (b) removing unbound material from the tissue protective cytokine receptor complex; and
- 20 (c) detecting the level of the first and second labels wherein if the second label is present the compound binds the complex and if the level of the first label decreases relative to the level of the first label where the labeled ligand is contacted with a tissue protective cytokine receptor complex under conditions conducive to binding in the absence of a test compound after removal of unbound material,
- 25

then a compound that binds to a tissue protective cytokine receptor complex is identified.

23. A method for identifying a compound that modulates the binding of a tissue protective cytokine receptor complex ligand to a tissue protective cytokine receptor complex, comprising:

30

(a) contacting a tissue protective cytokine receptor complex ligand with a tissue protective cytokine receptor complex in the presence of one or more test compounds under conditions conducive to binding; and

(b) measuring the amount of tissue protective cytokine receptor complex ligand bound to the tissue protective cytokine receptor complex;

such that if the amount of bound tissue protective cytokine receptor complex ligand measured in (b) differs from the amount of bound tissue protective cytokine receptor complex ligand measured in the absence of the one or more test compounds, then a compound that modulates the binding of a tissue protective cytokine receptor complex ligand to the tissue protective cytokine receptor complex is identified.

24. The method of Claim 23, wherein the amount of bound tissue protective cytokine receptor complex ligand is measured using a tissue protective cytokine receptor complex ligand-specific antibody.

25. The method of Claim 23, wherein the tissue protective cytokine receptor complex ligand is labeled and binding of the tissue protective cytokine receptor complex ligand to the tissue protective cytokine receptor complex is measured by detecting the label attached to the tissue protective cytokine receptor complex ligand.

26. The method of any one of Claim 23, wherein the tissue protective cytokine receptor complex ligand is labeled and binding of the labeled ligand to the tissue protective cytokine receptor complex is measured by detecting the label attached to the tissue protective cytokine receptor complex ligand.

27. The method of Claim 24, wherein the label is fluorescent.

28. A method for identifying a compound that modulates the interaction between a tissue protective cytokine receptor complex and a tissue protective cytokine receptor complex ligand, comprising:

(a) contacting a tissue protective cytokine receptor complex with one or more test compounds; and

(b) measuring the tissue protective cytokine receptor complex activity,

such that if the activity measured in (b) differs from the tissue protective cytokine receptor complex activity in the absence of the one or more test compounds, then a compound that modulates the interaction between the tissue protective cytokine receptor complex and the tissue protective cytokine receptor complex ligand is identified.

5     29.     The method of any one of Claims 1, 5, 11, 12, or 28, wherein the tissue protective cytokine receptor complex activity is measured by cell proliferation or cell differentiation.

10     30.     The method of any one of Claims 1, 5, 11, 12, or 28, wherein the tissue protective cytokine receptor complex activity measured is the ability of the tissue protective cytokine receptor complex to interact with a tissue protective cytokine receptor complex ligand.

31.     The method of any one of Claims 1, 5, 11, 12, 13, 22, or 28, wherein the step of assaying the identified compound for tissue protective activity comprises detecting the presence of nucleolin in the cell.

15     32.     The method of any one of Claims 1, 5, 11, 12, 13, 22, or 28, wherein the step of assaying the identified compound for tissue protective activity comprises detecting or measuring an increased level of activity of neuroglobin or cytoglobin in a cell.

33.     The method of any one of Claims 1, 22, 23, or 28, wherein the tissue protective cytokine receptor complex is in solution.

20     34.     The method of any one of Claims 1, 22, 23, or 28, wherein the tissue protective cytokine receptor complex is in a cell.

35.     The method of Claim 23, or 28, wherein the compound inhibits the binding of a tissue protective cytokine receptor complex ligand to a tissue protective cytokine receptor complex.

25     36.     The method of Claim 23, or 28, wherein the compound enhances the binding of a tissue protective cytokine receptor complex ligand to a tissue protective cytokine receptor complex.



37. The method of any one of Claims 1, 22, 23, or 28, wherein the tissue protective cytokine receptor complex contacted in step (a) is on a cell surface.
38. The method of any one of Claims 1, 22, 23, or 28, wherein the tissue protective cytokine receptor complex is on an isolated cell membrane.
- 5 39. The method of any one of Claims 1, 22, 23, or 28, wherein the tissue protective cytokine receptor complex is immobilized to a solid surface.
40. The method of Claim 39, wherein the solid surface is a microtiter dish.
41. The method of Claim 39, wherein the solid surface is a chip.
42. A method for identifying a compound that binds a tissue protective cytokine  
10 receptor complex, comprising:
- (a) contacting a test compound with a ligand-binding tissue protective receptor complex fragment comprising at least one EPO receptor extracellular domain and at least one  $\beta$  common receptor extracellular domain fused to an F<sub>c</sub> fragment attached to a solid support; and
  - 15 (b) removing unbound test compounds from the solid support;
  - (c) identifying the compound attached to the tissue protective cytokine receptor complex fragment,
- such that a compound bound to the solid support is identified as a compound that binds to a tissue protective cytokine receptor complex.
- 20 43. The method of any one of Claims 1, 5, 11, 12, 13, 15, 16, 22, 23, 28, or 42, wherein the test compound is an antibody specific for the tissue protective cytokine receptor complex.
44. The method of any one of Claims 16, 22, 23, 28, or 42, wherein the test  
25 compound is an antibody is specific for a tissue protective cytokine receptor complex ligand.
45. The method of any one of Claims 1, 5, 11, 12, 13, 15, 16, 22, 23, 28, or 42, wherein the test compound is a small molecule.

46. The method of any one of Claims 1, 5, 11, 12, 13, 15, 16, 22, 23, 28, or 42, wherein the test compound is a peptide.

47. The method of any one of Claims 1, 5, 11, 12, 13, 15, 16, 22, 23, 28, or 42, wherein the test compound is a member of a library.

5 48. The method of any one of Claims 16, 22, 23, 28, or 42, wherein the tissue protective cytokine receptor complex ligand is an EPO.

49. The method of any one of Claims 1, 5, 11, 12, 13, 28, or 42, wherein the compound binds the tissue protective cytokine receptor complex.

10 50. The method of any one of Claims 16, 23, 28, or 42, wherein the compound binds the tissue protective cytokine receptor complex ligand.

51. A method for identifying a compound that modulates a tissue protective activity in a mammal, comprising:

- 15 (a) administering the compound to a first animal immediately following infliction of an injury, wherein the first animal endogenously expresses a tissue protective cytokine receptor complex; and
- (b) administering the compound to a second animal immediately following infliction of the same injury as in step (a), wherein the second animal is deficient in expression of a tissue protective cytokine receptor complex;
- 20

such that if recovery from the injury differs in the animal of step (a) as compared to the animal of step (b), a compound that modulates a tissue protective activity is identified.

52. A method of treating or preventing a disease or disorder in a human comprising administering a therapeutically effective amount of a compound that modulates the activity of a tissue protective cytokine receptor complex to a human in need of such treatment or prevention, with the proviso that the compound is not an EPO.

25

53. The method of claim 52 wherein the disease or disorder is a disease or disorder of or affecting excitable tissue.

54. A method for enhancing the function of excitable cells, tissues, or organs in a human comprising administering a therapeutically effective amount of a compound that  
5 modulates the activity of a tissue protective cytokine receptor complex to a human in need thereof, wherein said enhancing the function of excitable cells, tissues, or organs results in the enhancement of associative learning or memory, with the proviso that the compound is not an EPO.

55. A method for modulating the activity of a tissue protective cytokine receptor  
10 complex in a human, comprising administering a compound that modulates the activity of said complex to a human in need of such treatment or prevention, with the proviso that the compound is not an EPO.

56. A method for treating or preventing a disease or disorder in a human comprising modulating a tissue protective cytokine receptor complex in a human.

15 57. A method for enhancing the function of excitable cells, tissues, or organs in a human comprising modulating a tissue protective cytokine receptor complex in a human.

58. The method of any one of claims 52, 53, 54, or 55 wherein the compound is identified by the method of any one of claims 1-5, 11-15, 21-23, 28, and 42.

59. The method of any one of claims 52, 53, 54, or 55 wherein the compound is an  
20 antibody specific for the tissue protective cytokine receptor complex.

60. The method of any one of claims 52, 53, 54, or 55 wherein the compound is an antibody is specific for a tissue protective cytokine receptor complex ligand.

61. The method of any one of claims 52, 53, 54, or 55 wherein the compound is a small molecule, peptide, or a member of a library.

25 62. The method of any one of claims 52, 53, 54, or 55 wherein the compound binds to the tissue protective cytokine receptor complex.

63. The method of any one of claims 52, 53, 54, or 55 wherein the compound enhances the activity of the tissue protective cytokine receptor complex.
64. The method of any one of claims 52, 53, 54, or 55 wherein the compound interferes with the activity of the tissue protective cytokine receptor complex.
- 5 65. The method of any one of claims 52, 53, 54, or 55 wherein the compound is administered in conjunction with an EPO.
66. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is a mood disorder, an anxiety disorder, depression, autism, an attention deficit hyperactivity disorder, Alzheimer's disease, aging, or cognitive dysfunction.
- 10 67. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is caused by hypoxia, seizure disorders, neurodegenerative diseases, neurotoxin poisoning, multiple sclerosis, hypotension, cardiac arrest, radiation, or hypoglycemia.
68. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is ischemia or stroke.
- 15 69. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is a neurodegenerative condition.
70. The method of claim 69 wherein the neurodegenerative condition is stroke, Alzheimer's disease, Parkinson's disease, cerebral palsy, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism, aging, or neuronal loss.
- 20 71. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is a neuromuscular or muscular condition.
72. The method of claim 71 wherein the neuromuscular or muscular condition is of a heart tissue.

73. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is of the heart.

74. The method of claim 73 wherein the disease or disorder is congestive heart failure, coronary artery disease, ischemia, myocardial infarction, angina, congenital heart disease, Prinzmetal angina, cardiac rupture, angiitis, arrhythmia, tachy-bradyarrhythmia, supraventricular, ventricular, conduction abnormalities, myocarditis, cor pulmonale, or Blunt and penetrating trauma.

75. The method of any one of claims 52, 53, or 56 wherein the disease or disorder is of the eye.

76. The method of claim 73 wherein the disease or disorder is optic neuritis, blunt and penetrating injuries, infections, sarcoid, sickle cell disease, retinal detachment, temporal arteritis, retinal ischemia, macular degeneration, retinitis pigmentosa, arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, diabetic retinopathy, or macular edema.

77. The method of claim 54 or 57 wherein the excitable tissue is central nervous system tissue or peripheral nervous system tissue.

78. The method of claim 77 wherein the peripheral nervous system tissue is placental tissue, retinal tissue, or cardiac tissue.

79. The method of any one of claims 52, 53, 54, or 55 wherein the administration is oral, topical, intracranial, intraluminal, inhalation, parenteral, intravenous, peripherally, intraarterial, subcutaneous, intramuscular, intraperitoneal, submucosal or intradermal administration.

80. The method of any one of claims 52, 53, 54, or 55 wherein the administration is prior to a medical, surgical, or natural procedure.

81. The method of claim 80 wherein the surgical procedure is tumor resection, aneurysm repair, or a coronary artery bypass procedure.

82. The method of claim 80 wherein the medical procedure is chemotherapy.
83. The method of claim 80 wherein the natural procedure is labor or delivery.




FIG.1

2/15




FIG.2



3/15




FIG.3

SUBSTITUTE SHEET (RULE 26)

4/15




FIG. 1

5/15




FIG.5

6/15




FIG.6

7/15




FIG.7

8/15




FIG.8

9/15




FIG.9

10/15




FIG.10



11/15


\*  $p = 0.05$ 

FIG. 11A

12/15




FIG. 11B

13/15




FIG.12

14/15




FIG.13

15/15




FIG.14

## SEQUENCE LISTING

<110> The Kenneth Warren Institute, Inc.

<120> TISSUE PROTECTIVE CYTOKINE RECEPTOR COMPLEX, ASSAYS FOR IDENTIFYING  
TISSUE PROTECTIVE COMPOUNDS AND USES THEREOF

<130> 10165-028-228

<140>  
<141>

<150> 10/676,694  
<151> 2003-09-30

<150> 60/465,891  
<151> 2003-04-25

<160> 212

<170> PatentIn version 3.2

<210> 1  
<211> 5  
<212> PRT  
<213> Homo sapiens

<400> 1

Val Leu Gln Arg Tyr  
1 5

<210> 2  
<211> 8  
<212> PRT  
<213> Homo sapiens

<400> 2

Thr Lys Val Asn Phe Tyr Ala Trp  
1 5

<210> 3  
<211> 9  
<212> PRT  
<213> Homo sapiens

<400> 3

Ser Gly Leu Arg Ser Leu Thr Thr Leu  
1 5

<210> 4  
<211> 6  
<212> PRT  
<213> Homo sapiens

<400> 4

Ser Asn Phe Leu Arg Gly  
1 5

<210> 5

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 5

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 6  
 <211> 193  
 <212> PRT  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 6

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp



180

185

190

Arg

<210> 7  
 <211> 580  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 atgggggtgc acgaatgtcc tgccctggctg tggcttctcc tgtccctgct gtcgctccct 60  
 ctgggcctcc cagtcctggg cgcctcatct gtgacagccg agtcctggag 120  
 aggtacctct tggaggccaa ggaggccgag aatatcacga cgggctgtgc tgaacactgc 180  
 agcttgaatg agaatatcac tgtcccagac accaaagtta atttctatgc ctggaagagg 240  
 atggagggtcg ggcagcaggc cgtagaagtc tggcagggcc tggccctgct gtcggaagct 300  
 gtccctgcggg gccaggccct gttggtcaac tcttccagc cgtgggagcc cctgcactgc 360  
 atgtggataa agccgtcagt ggccttcgca gcctcaccac tctgcttcgg gctctgggag 420  
 cccagaagga agccatctcc cctccagatg cggcctcagc tgctccactc cgaacaatca 480  
 ctgctgacac ttctcgaaac tcttccgagt ctactccaat ttcctccggg gaaagctgaa 540  
 gctgtacaca ggggaggcct gcaggacagg ggacagatga 580

<210> 8  
 <211> 35  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 8  
 agctctcgag gcgcggagat ggggggtgcac gaatg 35

<210> 9  
 <211> 36  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 9  
 atgctctaga cacacctggt catctgtccc ctgtcc 36

<210> 10  
 <211> 193  
 <212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 11

&lt;211&gt; 45

&lt;212&gt; DNA

&lt;213&gt; Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 11  
 catgtggata aagccgtcga gggccttcgc agcctcacca ctctg 45

<210> 12  
 <211> 45  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 12  
 cagagtgggtg aggctgacga ggccttcgac ggctttatcc acatg 45

<210> 13  
 <211> 45  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 13  
 gagaatatca ctgtcccaga caccgacgtt aattttctatg cctgg 45

<210> 14  
 <211> 45  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: primer

<400> 14  
 ccaggcatag aaattaacgt cgggtgtctgg gacagtgata ttctc 45

<210> 15  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 15

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ala Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 16  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 16

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Ala Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 17  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 17

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu

20

25

30

Ile Cys Asp Ser Ile Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 18

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 18

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
           20                          25                          30  
 Ile Cys Asp Ser Arg Ser Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
           35                          40                          45  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
           50                          55                          60  
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
           65                          70                          75                          80  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
                           85                          90                          95  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
                           100                          105                          110  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
                           115                          120                          125  
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
           130                          135                          140  
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
           145                          150                          155                          160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
                           165                          170                          175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
                           180                          185                          190

Arg

<210> 19  
 <211> 193  
 <212> PRT  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: mutein  
  
 <400> 19

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1                          5                          10                          15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
                   20                                  25                                  30  
 Ile Cys Asp Ser Arg Val Ala Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
           35                                  40                                  45  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
           50                                  55                                  60  
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
   65                                  70                                  75                                  80  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
                   85                                  90                                  95  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
                   100                                  105                                  110  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
           115                                  120                                  125  
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
   130                                  135                                  140  
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
  145                                  150                                  155                                  160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
           165                                  170                                  175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
           180                                  185                                  190

Arg

<210> 20  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 20

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu



```

1           5           10           15
Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
      20           25           30
Ile Cys Asp Ser Arg Val Leu Ala Arg Tyr Leu Leu Glu Ala Lys Glu
      35           40           45
Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
      50           55           60
Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
      65           70           75           80
Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
      85           90           95
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
      100          105          110
Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
      115          120          125
Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
      130          135          140
Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
      145          150          155          160
Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
      165          170          175
Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
      180          185          190

```

Arg

```

<210> 21
<211> 193
<212> PRT
<213> Artificial

<220>
<223> Description of Artificial Sequence: mutein

<400> 21

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15  
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30  
 Ile Cys Asp Ser Arg Val Leu Glu Ala Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60  
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125  
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140  
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 22  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

&lt;400&gt; 22

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 23

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 23

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Glu Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 24

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

<223> Description of Artificial Sequence: mutein

<400> 24

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Gln Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 25

<211> 193

<212> PRT

<213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 25

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ala Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 26

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 26

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Phe Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 27

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 27

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ile Leu Leu Glu Ala Glu Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 28



<211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 28

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Glu Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 29  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 29

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Ala Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 30  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 30

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Ala  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 31  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 31

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 32

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 32

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Ser Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 33  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 33

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1              5              10              15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
              20              25              30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
      35              40              45

Ala Glu Asn Ile Thr Thr Gly Tyr Ala Glu His Cys Ser Leu Asn Glu
      50              55              60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65              70              75              80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
              85              90              95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
              100              105              110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
              115              120              125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
              130              135              140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile
145              150              155              160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
              165              170              175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
              180              185              190

```

Arg

<210> 34  
 <211> 193  
 <212> PRT  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 34

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Asn Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp

180

185

190

Arg

&lt;210&gt; 35

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 35

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu Thr Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175



Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 36  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 36

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Ser Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 37  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 37

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Tyr Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu

165

170

175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 38  
 <211> 193  
 <212> PRT  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 38

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Lys Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 39  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein  
 <400> 39

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Lys Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 40  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 40

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Asn Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 41  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 41

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Ala Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile

145                                      150                                      155                                      160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
    165                                      170                                      175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
    180                                      185                                      190  
 Arg  
  
 <210> 42  
 <211> 193  
 <212> PRT  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: mutein  
  
 <400> 42  
  
 Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1                                      5                                      10                                      15  
  
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
    20                                      25                                      30  
  
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
    35                                      40                                      45  
  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
    50                                      55                                      60  
  
 Asn Ile Thr Val Pro Ala Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65                                      70                                      75                                      80  
  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
    85                                      90                                      95  
  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
    100                                      105                                      110  
  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
    115                                      120                                      125  
  
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
    130                                      135                                      140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 43  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 43

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Ile Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140



Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg )

<210> 44

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 44

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu

130

135

140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 45

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 45

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 46  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 46

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Ala Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 47  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 47

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Ala Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 48  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 48

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Ile Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly

115

120

125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 49  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 49

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Ala Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 50  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 50

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Ala Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 51

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 51

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Ser Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser



100

105

110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 52  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 52

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Phe Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 53

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 53

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Asn Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 54  
 <211> 193  
 <212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 54

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Ala Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 55

<211> 193

<212> PRT

<213> Artificial .

<220>

<223> Description of Artificial Sequence: mutein

<400> 55

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Asn Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
                     85                    90                    95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
                     100                    105                    110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
                     115                    120                    125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
                     130                    135                    140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
                     145                    150                    155                    160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
                     165                    170                    175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
                     180                    185                    190

Arg

<210> 56  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 56

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
   1                    5                    10                    15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
                     20                    25                    30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
                     35                    40                    45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
                     50                    55                    60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
   65                    70                    75                    80

Met Glu Val Gly Gln Gln Ala Val Thr Val Trp Gln Gly Leu Ala Leu  
                   85                                  90                                  95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
                   100                                  105                                  110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
                   115                                  120                                  125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
                   130                                  135                                  140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
                   145                                  150                                  155                                  160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
                   165                                  170                                  175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
                   180                                  185                                  190

Arg

<210> 57

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 57

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
   1                  5                                  10                                  15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
                   20                                  25                                  30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
                   35                                  40                                  45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
                   50                                  55                                  60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg

Arg

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu
1				5					10					15	
Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		
Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu
		35					40					45			
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu
	50					55					60				

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Ala Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 59  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 59

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60



Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Arg Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 60  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 60

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 61  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 61

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu

50

55

60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Arg Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 62  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 62

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 63  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 63

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ala Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 64

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 64

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu

35

40

45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Thr Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 65  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 65

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Ala  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 66

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 66

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Ile  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 67

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 67

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30



Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Ala Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 68  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 68

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu

20

25

30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Ala Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 69

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 69

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30  
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60  
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125  
 Leu Glu Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140  
 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 70

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 70

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ala Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala<sup>1</sup>Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 71  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 71

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu

Arg

64

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Ala Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 73  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 73

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Ala Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 74  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

&lt;400&gt; 74

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Ile Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 75

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein



&lt;400&gt; 75

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Ala Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 76

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

<223> Description of Artificial Sequence: mutein

<400> 76

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Leu Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 77

<211> 193

<212> PRT

<213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 77

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Lys Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 78

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 78

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Ala Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 79

&lt;211&gt; 193

<212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 79

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Ser Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 80

<211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 80

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Ala Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 81  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 81

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1           5           10           15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
          20           25           30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
          35           40           45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
          50           55           60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65           70           75           80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85           90           95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
          100          105          110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
          115          120          125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
          130          135          140

Ala Ile Ser Pro Pro Asp Ala Ala Ala Ala Ala Pro Leu Arg Thr Ile
145           150           155           160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
          165          170          175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180          185          190

```

Arg

<210> 82  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 82

```

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
1           5           10           15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
          20           25           30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu
      35           40           45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu
      50           55           60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg
65           70           75           80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu
          85           90           95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser
      100           105           110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly
      115           120           125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu
      130           135           140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Ala Ile
145           150           155           160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu
          165           170           175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp
          180           185           190

```

Arg



<210> 83  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 83

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ala  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 84  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 84

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Ala Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

&lt;210&gt; 85

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 85

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 86  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 86

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Ile Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp

180

185

190

Arg

&lt;210&gt; 87

&lt;211&gt; 193

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 87

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Ala Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 88  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 88

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ala Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 89  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 89

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Lys Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 90  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 90

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160



Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Ala Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 91  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 91

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Tyr Leu  
                   165                                  170                                  175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
                   180                                  185                                  190

Arg

<210> 92  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 92

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1                  5                                  10                                  15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
                   20                                  25                                  30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
                   35                                  40                                  45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
                   50                                  55                                  60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65                                  70                                  75                                  80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
                   85                                  90                                  95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
                   100                                  105                                  110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
                   115                                  120                                  125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
                   130                                  135                                  140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145                                  150                                  155                                  160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Ala Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 93  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 93

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile

145                                      150                                      155                                      160  
 Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Ala  
    165                                      170                                      175  
 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
    180                                      185                                      190  
  
 Arg  
  
 <210> 94  
 <211> 193  
 <212> PRT  
 <213> Artificial  
  
 <220>  
 <223> Description of Artificial Sequence: mutein  
  
 <400> 94  
  
 Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1                                      5                                      10                                      15  
  
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
    20                                      25                                      30  
  
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
    35                                      40                                      45  
  
 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
    50                                      55                                      60  
  
 Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65                                      70                                      75                                      80  
  
 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
    85                                      90                                      95  
  
 Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
    100                                      105                                      110  
  
 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
    115                                      120                                      125  
  
 Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
    130                                      135                                      140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Ala Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 95  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 95

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Glu Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 96  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 96

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu

130

135

140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Ala Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 97  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein  
 <400> 97

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 98  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 98

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125



Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Trp Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 99  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 99

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Ala Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 100  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 100

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Ala Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg .

<210> 101  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein  
 <400> 101

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Ala Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 102  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 102

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Ala Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 103  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 103

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Ser Arg Thr Gly Asp  
 180 185 190

Arg

<210> 104  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 104

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Ala Arg Thr Gly Asp  
 180 185 190

Arg

<210> 105  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 105

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu

85

90

95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Ala Thr Gly Asp  
 180 185 190

Arg

<210> 106

<211> 192

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 106

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Glu His Cys Ser Leu Asn Glu Asn  
 50 55 60

Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg Met  
 65 70 75 80



Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu Leu  
                             85                            90                            95

Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser Gln  
                             100                            105                            110

Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Glu Gly Leu  
                             115                            120                            125

Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu Ala  
                             130                            135                            140

Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile Thr  
                             145                            150                            155                            160

Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu Arg  
                             165                            170                            175

Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp Arg  
                             180                            185                            190

<210> 107  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 107

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1                            5                            10                            15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
                             20                            25                            30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
                             35                            40                            45

Ala Glu Asn Ile Thr Thr Gly Cys Asn Glu Thr Cys Ser Leu Asn Glu  
                             50                            55                            60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65                            70                            75                            80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
                             85                            90                            95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 108

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 108

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Asp Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80



Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Glu Ser Leu Thr Thr Ser Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 110  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 110

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Ala Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 111  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 111

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg

65		70		75		80									
Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu
			85					90					95		
Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser
			100				105						110		
Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly
		115				120						125			
Leu	Arg	Ser	Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu
	130					135					140				
Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile
145					150				155						160
Thr	Ala	Asp	Thr	Phe	Arg	Ala	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu
				165					170					175	
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp
			180					185					190		

Arg

<210> 112  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein  
 <400> 112

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu
1				5					10					15	
Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu
			20					25					30		
Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu
		35					40					45			
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu
	50					55					60				

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 113  
 <211> 193  
 <212> PRT  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: mutein

<400> 113

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 114  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 114

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu



50

55

60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 115  
 <211> 193  
 <212> PRT  
 <213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: mutein

&lt;400&gt; 115

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Ala Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 116  
<211> 193  
<212> PRT  
<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 116

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Asn Ile Thr Val Pro Asp Thr Ala Val Asn Phe Tyr Ala Trp Ala Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Ala Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Ala Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Ala Leu Ala Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 117  
<211> 193  
<212> PRT  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: mutein

<400> 117

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60

Lys Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Lys Ser Ser  
100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190

Arg

<210> 118

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 118

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Arg Ala Leu Leu Glu Ala Lys Glu

35

40

45

Ala Glu Lys Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 119

<211> 193

<212> PRT

<213> Artificial

<220>

<223> Description of Artificial Sequence: mutein

<400> 119

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Ser Leu  
 1 5 10 15

Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
 20 25 30

Ile Cys Asp Ser Arg Val Leu Glu Ala Ala Leu Leu Glu Ala Lys Glu  
 35 40 45

Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
 50 55 60

Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
 65 70 75 80

Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
 85 90 95

Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
 100 105 110

Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
 115 120 125

Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
 130 135 140

Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
 145 150 155 160

Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
 165 170 175

Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
 180 185 190

Arg

<210> 120

<211> 36

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 120

gtctactcca atttcctcga gggaaagctg aagctg

36

<210> 121

<211> 34

<212> DNA

<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 121  
gcttcagctt tccctcgagg aaattggagt agac 34

<210> 122  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 122  
ccgtcagtg ccttgagagc ctcaccactc tg 32

<210> 123  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 123  
cagagtggcg aggctctcaa ggccactgac gg 32

<210> 124  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 124  
ccgtcagtg ccttgagagc ctcaccactc tg 32

<210> 125  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 125  
cagagtggcg aggctctcaa ggccactgac gg 32

<210> 126  
<211> 31  
<212> DNA  
<213> Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 126

cgcagcctca ccacttcgct tcgggctctg g

31

&lt;210&gt; 127

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 127

ccagagcccg aagcgaagtg gtgaggctgc g

31

&lt;210&gt; 128

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 128

gaatatcact gtcccagacg gtggtgcctg gaagaggatg

40

&lt;210&gt; 129

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 129

catcctcttc caggcaccac cgtctgggac agtgatattc

40

&lt;210&gt; 130

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer

&lt;400&gt; 130

tacctcttgg aggccgcgga ggccgagaat atc

33

&lt;210&gt; 131

&lt;211&gt; 33

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;



<223> Description of Artificial Sequence: primer

<400> 131

gatattctcg gcctccgcgg cctccaagag gta

33

<210> 132

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 132

gctgacactt tccgcgcact cttccgagtc tactc

35

<210> 133

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 133

gagtagactc ggaagagtgc gcggaaagtg tcagc

35

<210> 134

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 134

atttcctccg gggagcgctg aagctgtaca cag

33

<210> 135

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 135

ctgtgtacag cttcagcgct ccccggagga aat

33

<210> 136

<211> 32

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 136  
ctccggggaa agctggcgct gtacacaggg ga 32

<210> 137  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 137  
tcccctgtgt acagcgccag ctttccccgg ag 32

<210> 138  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 138  
actgtcccag acaccgcagt taatttctat gcctg 35

<210> 139  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 139  
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 140  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 140  
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 141  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 141  
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 142  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 142  
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 143  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 143  
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 144  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 144  
ctctgggagc ccaggcggaa gccatctccc ct 32

<210> 145  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 145  
aggggagatg gcttccgcct gggctcccag ag 32

<210> 146  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 146

gctgacactt tccgcgcact cttccgagtc tactc

35

<210> 147  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 147  
gagtagactc ggaagagtgc gcggaaagtg tcagc

35

<210> 148  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 148  
agttaatttc tatgcctggg cgaggatgga ggtcg

35

<210> 149  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 149  
cgacctccat cctcgcccag gcatagaaat taact

35

<210> 150  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 150  
gctgacactt tccgcgcact cttccgagtc tactc

35

<210> 151  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 151  
gagtagactc ggaagagtgc gcggaaagtg tcagc

35

<210> 152  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 152  
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 153  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 153  
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 154  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 154  
actgtcccag acaccgcagt taatttctat gcctg 35

<210> 155  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 155  
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 156  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 156  
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 157  
<211> 33  
<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 157  
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 158  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 158  
atttcctccg gggagcgctg aagctgtaca cag 33

<210> 159  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 159  
ctgtgtacag cttcagcgct ccccgaggga aat 33

<210> 160  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 160  
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 161  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 161  
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 162  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 162  
atttcctccg gggagcgctg aagctgtaca cag

33

<210> 163  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 163  
ctgtgtacag cttcagcgct ccccggagga aat

33

<210> 164  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 164  
actgtcccag acaccgcagt taatttctat gcctg

35

<210> 165  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 165  
cagggcataga aattaactgc ggtgtctggg acagt

35

<210> 166  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 166  
tgcagctgca tgtggatgca gccgtcagtg gcc

33

<210> 167  
<211> 33  
  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Description of Artificial Sequence: primer  
  
<400> 167  
ggccactgac ggctgcatcc acatgcagct gca 33  
  
<210> 168  
<211> 33  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Description of Artificial Sequence: primer  
  
<400> 168  
atttcctccg gggagcgcgtg aagctgtaca cag 33  
  
<210> 169  
<211> 33  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Description of Artificial Sequence: primer  
  
<400> 169  
ctgtgtacag cttcagcgcgt ccccgaggga aat 33  
  
<210> 170  
<211> 35  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Description of Artificial Sequence: primer  
  
<400> 170  
actgtcccag acaccgcagt taatttctat gcctg 35  
  
<210> 171  
<211> 35  
<212> DNA  
<213> Artificial  
  
<220>  
<223> Description of Artificial Sequence: primer  
  
<400> 171  
caggcataga aattaactgc ggtgtctggg acagt 35



<210> 172  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 172  
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 173  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 173  
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 174  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 174  
tgcagctgca tgtggatgca gccgtcagtg gcc 33

<210> 175  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 175  
ggccactgac ggctgcatcc acatgcagct gca 33

<210> 176  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 176  
atttcctccg gggagcgctg aagctgtaca cag 33

<210> 177

<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 177  
ctgtgtacag cttcagcgct ccccgaggga aat 33

<210> 178  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 178  
actgtcccag acaccgcagt taatttctat gcctg 35

<210> 179  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 179  
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 180  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 180  
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 181  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 181  
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 182  
<211> 35

<212> DNA  
<213> Artificial  
<220>  
<223> Description of Artificial Sequence: primer  
<400> 182  
gctgacactt tccgcgcact cttccgagtc tactc 35  
  
<210> 183  
<211> 35  
<212> DNA  
<213> Artificial  
<220>  
<223> Description of Artificial Sequence: primer  
<400> 183  
gagtagactc ggaagagtgc gcggaaagtg tcagc 35  
  
<210> 184  
<211> 33  
<212> DNA  
<213> Artificial  
<220>  
<223> Description of Artificial Sequence: primer  
<400> 184  
tgcagctgca tgtggatgca gccgtcagtg gcc 33  
  
<210> 185  
<211> 33  
<212> DNA  
<213> Artificial  
<220>  
<223> Description of Artificial Sequence: primer  
<400> 185  
ggccactgac ggctgcatcc acatgcagct gca 33  
  
<210> 186  
<211> 33  
<212> DNA  
<213> Artificial  
<220>  
<223> Description of Artificial Sequence: primer  
<400> 186  
atttcctccg gggagcgctg aagctgtaca cag 33  
  
<210> 187  
<211> 33  
<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 187  
ctgtgtacag cttcagcgct ccccgaggga aat 33

<210> 188  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 188  
actgtcccag acaccgcagt taatttctat gcctg 35

<210> 189  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 189  
caggcataga aattaactgc ggtgtctggg acagt 35

<210> 190  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 190  
agttaatttc tatgcctggg cgaggatgga ggtcg 35

<210> 191  
<211> 35  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 191  
cgacctccat cctcgcccag gcatagaaat taact 35

<210> 192  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 192  
gctgacactt tccgcgcact cttccgagtc tactc 35

<210> 193  
<211> 35  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 193  
gagtagactc ggaagagtgc gcggaaagtg tcagc 35

<210> 194  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 194  
ctccggggag cgctggcgct gtacacaggg ga 32

<210> 195  
<211> 32  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 195  
tcccctgtgt acagcgccag cgctccccgg ag 32

<210> 196  
<211> 31  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 196  
caaggaggcc gagaaaatca cgacgggctg t 31

<210> 197  
<211> 31  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 197  
acagcccgtc gtgattttct cggcctcctt g 31

<210> 198  
<211> 37  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 198  
actgcagctt gaatgagaaa atcactgtcc cagacac 37

<210> 199  
<211> 37  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 199  
gtgtctggga cagtgatttt ctcattcaag ctgcagt 37

<210> 200  
<211> 31  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 200  
aggccctggt ggtcaaactt tcccagccgt g 31

<210> 201  
<211> 31  
<212> DNA  
<213> Artificial

<220>  
<223> Description of Artificial Sequence: primer

<400> 201  
cacggctggg aagatttgac caacagggcc t 31

<210> 202  
<211> 33  
<212> DNA  
<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 202

atttcctccg gggatggctg aagctgtaca cag

33

<210> 203

<211> 33

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 203

ctgtgtacag cttcagccat ccccggagga aat

33

<210> 204

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 204

agccgagtc tggaggcggc cctcttgag gccaa

35

<210> 205

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 205

ttggcctcca agagggcgc ctccaggact cggct

35

<210> 206

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 206

agccgagtc tggagagggc cctcttgag gccaa

35

<210> 207

<211> 35

<212> DNA

<213> Artificial

<220>

<223> Description of Artificial Sequence: primer

<400> 207  
 ttggcctcca agagggccct ctccaggact cggct 35

<210> 208  
 <211> 6059

<212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: plasmid

<400> 208  
 ctagagtcga cccgggcggc cgcttcctt tagtgagggt taatgcttcg agcagacatg 60  
 ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120  
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180  
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240  
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300  
 ggcgtaatat cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360  
 gcgaatggac ggcgccgtga gcggcgcat aagcgcggcg ggtgtggtgg ttacgcgcag 420  
 cgtgaccgct acacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt 480  
 tctcgccacg ttcgccggct tccccgtca agctctaaat cgggggctcc ctttaggggtt 540  
 ccgatttagt gctttacggc acctcgacc caaaaaactt gattaggggtg atggttcacg 600  
 tagtgggcca tcgccctgat agacggtttt tcgccctttg acgttggagt ccacgttctt 660  
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt 720  
 tgatttataa gggattttgc cgatttcggc ctattgggtta aaaaatgagc tgatttaaca 780  
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtathtt 840  
 ctccctacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900  
 ctgaaataac ctctgaaaga ggaacttgggt taggtacctt ctgaggcgga aagaaccagc 960  
 tgtggaatgt gtgtcagtta ggggtgtgaa agtccccagg ctccccagca ggcagaagta 1020  
 tgcaaagcat gcactcfaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080  
 caggcagaag tatgcaaagc atgcactcfa attagtcagc aaccatagtc ccgcccctaa 1140  
 ctccgcccct cccgccccta actccgccc gttccgccc ttctccgccc catggctgac 1200  
 taattttttt tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260  
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320  
 cagtctcgaa ctttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtt 1380



tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg	1440
ctctgatgcc gccgtgttcc ggctgtcagc gcaggggccc cgggttcttt ttgtcaagac	1500
cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc	1560
cacgacgggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg	1620
gctgctattg ggcgaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga	1680
gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg	1740
cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaagccgg	1800
tcttgctgat caggatgatc tggacgaaga gcatcagggg ctgcgccag ccgaactgtt	1860
cgccaggctc aaggcgcgca tgcccgacgg cgaggatctc gtcgtgacct atggcgatgc	1920
ctgcttgccg aatatcatgg tggaatatgg ccgcttttct ggattcatcg actgtggccg	1980
gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga	2040
gcttggcggc gaatgggctg accgcttctc cgtgctttac ggtatcgccg ctcccgattc	2100
gcagcgcata gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc	2160
gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt	2220
attttcatta catctgtgtg ttggtttttt gtgtgaatcg atagcgataa ggatccgcgt	2280
atggtgcaat ctcatgataa tctgctctga tgccgcatag ttaagccagc cccgacaccc	2340
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat	2520
ggttttcttag acgtcagggt gcacttttct gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcatttttgc ttctgttttt tgctcaccca gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gcccogaaga acgttttcca atgatgagca cttttaaagt	2880
tctgctatgt ggcgcggtat tatcccgat tgacgcgggg caagagcaac tcggtcgccg	2940
catacactat tctcagaatg acttgggttga gtactacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180

aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttccggct ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcactggggc cagatggtaa	3420
gccctcccg atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt	3540
ttactcatat atactttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt	3600
gaagatcctt ttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacia aaaaaccacc gctaccagcg gtggtttggt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccggggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttggagcg aacgacctac accgaactga gatacctaca	4080
gogtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccgg	4140
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta	4200
tctttatagt cctgtcgggt ttgccacct ctgacttgag cgtcgatttt tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ccttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgtat	4440
ctatatcata atatgtacat ttatatggc tcatgtccaa tatgaccgcc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggg cattagttca tagccatat	4560
atggagttcc gcgttacata acttacggta aatggcccgc ctggctgacc gcccaacgac	4620
ccccgccc atgacgtcaat aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtatttacgg taaactgcc acttggcagt acatcaagt	4740
tatcatatgc caagtccgcc ccctattgac gtcaatgacg gtaaatggcc cgctggcat	4800
tatgcccagt acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatgggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac	4980

```

caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc 5040
aaatgggagg taggcgtgta cggtagggagg tctatataag cagagctcgt ttagtgaacc 5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca 5160
gtgcttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc 5220
agaagttggg cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga 5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct 5340
attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat 5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga 5460
gatgggggtg cacgaatgtc ctgcctgggt gtggcttctc ctgtccctgc tgtcgctccc 5520
tctgggcttc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtcctgga 5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg 5640
cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag 5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcggaagc 5760
tgtcctgcgg ggccaggccc tgttgggtcaa ctcttccag ccgtgggagc ccctgcagct 5820
gcatgtggat aaagccgtca gtggccttcg cagcctcacc actctgcttc gggctctgcg 5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940
cactgctgac actttccgca aactcttcg agtctactcc aatttcctcc ggggaaagct 6000
gaagctgtac acaggggagg cctgcaggac aggggacat catcaccatc accattgat 6059

```

&lt;210&gt; 209

&lt;211&gt; 6059

&lt;212&gt; DNA

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: plasmid

&lt;400&gt; 209

```

ctagagtoga cccgggdcggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60
ataagatata ttgatgagtt tggacaaacc acaactagaa tgcagtgaag aaaatgcttt 120
atgtgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180
gttaacaaca acaattgcat tcatttttatg tttcagggtc agggggagat gtgggaggtt 240
ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300
ggcgtaatat cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360
gcgaatggac gcgccctgta gggcgccatt aagcgcgccg ggtgtggtgg ttacgcgcag 420
cgtgaccgct acacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt 480

```

tctcgccacg ttcgccgget ttccccgtca agctctaaat cgggggctcc ctttaggggtt	540
ccgatttagt gctttacggc acctcgaccc caaaaaactt gattaggggtg atgggttcacg	600
tagtggggcca tcgccctgat agacgggtttt tcgccctttg acgttggagt ccacgttctt	660
taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt	720
tgatttataa gggatttttg cgatttcggc ctattgggtta aaaaatgagc tgatttaaca	780
aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcgggtatttt	840
ctccttacgc atctgtgagg tattttcacac cgcatacgcg gatctgcgca gcaccatggc	900
ctgaaataac ctctgaaaga ggaacttggg taggtacctt ctgaggcgga aagaaccagc	960
tgtggaatgt gtgtcagtta ggggtgtggaa agtccccagg ctccccagca ggcagaagta	1020
tgcaaagcat gcattctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag	1080
caggcagaag tatgcaaagc atgcattctca attagtcagc aaccatagtc ccgcccctaa	1140
ctccgcccct cccgccccta actccgccc gttccgccc ttctccgccc catggctgac	1200
taattttttt tatttatgca gaggcggagg ccgcctcggc ctctgagcta ttccagaagt	1260
agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa	1320
cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtc	1380
tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg	1440
ctctgatgcc gcctgttcc ggctgtcagc gcagggggcg ccggttcttt ttgtcaagac	1500
cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc	1560
cacgacgggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg	1620
gctgctattg ggcgaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga	1680
gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg	1740
cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaaagccg	1800
tcttgctgat caggatgatc tggacgaaga gcatcagggg ctccgcccag ccgaactgtt	1860
cgccaggctc aaggcgcgca tgcccagcgg cgaggatctc gtcgtgaccc atggcgatgc	1920
ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg	1980
gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga	2040
gcttggcggc gaatgggctg accgcttctt cgtgctttac ggtatcgccg ctcccgatc	2100
gcagcgcata gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc	2160
gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt	2220
attttcatta catctgtgtg ttggtttttt gtgtgaatcg atagcgataa ggatccgcgt	2280

atggtgact ctcagtacaa tctgctctga tgccgcatag ttaagccagc cccgacaccc	2340
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat	2520
ggtttcttag acgtcagggtg gcacttttcg gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcattttgct ttccgtgttt tgctcaccca gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca cttttaaagt	2880
tctgctatgt ggcgcggtat tatcccgat tgacgccggg caagagcaac tcggtcgccg	2940
catacactat totcagaatg acttggttga gtactcacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttccggct ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gactggggc cagatggtaa	3420
gccctcccgat atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggttaactgt cagaccaagt	3540
ttactcatat atactttaga ttgatttaaa acttcatttt taatttataaa ggatctaggt	3600
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac ccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttggt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttgagcgc aacgacctac accgaactga gatacctaca	4080
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt	4140

aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta	4200
tctttatagt cctgtcgggt ttogccacct ctgacttgag cgtcgatttt tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ctttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgtat	4440
ctatatcata atatgtacat ttatatggc tcatgtccaa tatgaccgcc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggt cattagttca tagcccatat	4560
atggagttcc gcgttacata acttacggta aatggccgcg ctggctgacc gcccaacgac	4620
ccccgcccac tgacgtcaat aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtatttacgg taaactgcc acttggcagt acatcaagtg	4740
tatcatatgc caagtccgcc ccctattgac gtcaatgacg gtaaattggcc cgcctggcat	4800
tatgccagat acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac	4980
caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc	5040
aaatgggcgg taggcgtgta cggtgggagg tctatataag cagagctcgt ttagtgaacc	5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca	5160
gtgcttctga cacaacagtc togaacttaa gctgcagtga ctctcttaag gtagccttgc	5220
agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga	5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct	5340
attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat	5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga	5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgtctcc	5520
tctgggcctc ccagtcctgg gcgcccacc acgcctcatc tgtgacagcc gagtctgga	5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgta atgaaacctg	5640
cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag	5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc	5760
tgtcctgctg ggccaggccc tgttgggtcaa ctcttcccag ccgtgggagc ccctgcagct	5820
gcatgtggat aaagccgtca gtggccttcg cagcctcacc actctgcttc gggctctgctg	5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat	5940

cactgctgac actttccgca aactcttccg agtctactcc aatttcctcc ggggaaagct 6000  
 gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

<210> 210  
 <211> 6059  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: plasmid

<400> 210  
 ctagagtcga cccgggcggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60  
 ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120  
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180  
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240  
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300  
 ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360  
 gcgaatggac gcgccctgta gcggcgcatc aagcgcgccg ggtgtggtgg ttacgcgcag 420  
 cgtgaccgct aacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt 480  
 tctcgccacg ttcgccggct tccccgtca agctctaaat cgggggctcc ctttaggggt 540  
 ccgatttagt gctttacggc acctcgacct caaaaaactt gattaggggtg atggttcacg 600  
 tagtgggcca tcgccctgat agacggtttt tcgccctttg acgttggagt ccacgttctt 660  
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt 720  
 tgatttataa gggattttgc cgatttcggc ctattggtta aaaaatgagc tgatttaaca 780  
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtatttt 840  
 ctccctacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900  
 ctgaaataac ctctgaaaga ggaacttggt taggtacctt ctgaggcgga aagaaccagc 960  
 tgtggaatgt gtgtcagtta ggggtgtgaa agtccccagg ctccccagca ggcagaagta 1020  
 tgcaaagcat gcattctaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080  
 caggcagaag tatgcaaagc atgcattctc attagtcagc aaccatagtc ccgcccctaa 1140  
 ctccgcccac cccgccccta actccgcccga gttccgcccga ttctccgccc catggctgac 1200  
 taattttttt tattttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260  
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320  
 cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtc 1380  
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440

ctctgatgcc gccgtgttcc ggctgtcagc gcaggggagc ccggttcttt ttgtcaagac	1500
cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc	1560
cacgacgggc gttccttgcg cagctgtgct cgacgttgtc actgaagcgg gaagggactg	1620
gctgctattg ggccaagtgc cggggcagga tctcctgtca tctcaccttg ctctgccga	1680
gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg	1740
cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaagccgg	1800
tcttgtcgat caggatgatc tggacgaaga gcatcagggg ctccgcgccag ccgaactgtt	1860
cgccaggctc aaggcgcgca tgcccgacgg cgaggatctc gtcgtgaccc atggcgatgc	1920
ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg	1980
gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga	2040
gcttggcggc gaatgggctg accgcttctt cgtgctttac ggtatcgccg ctcccgattc	2100
gcagcgcac gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc	2160
gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt	2220
attttcatta catctgtgtg ttggtttttt gtgtgaatcg atagcgataa ggatccgcgt	2280
atggtgcact ctcagtacaa tctgctctga tgccgcatag ttaagccagc cccgacaccc	2340
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat	2520
ggtttcttag acgtcagggt gcacttttct gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcattttgct ttctgttttt tgctcaccca gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca cttttaaaagt	2880
tctgctatgt ggcgcggtat tatcccgat tgacgccggg caagagcaac tcggtcgccg	2940
catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaaa	3120
catgggggat catgtaaact gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240



aactggcgaa ctacttactc tagcttcccc gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttccggct ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcactggggc cagatggtaa	3420
gccctcccgat atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt	3540
ttactcatat atacttttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt	3600
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttggt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgtc ctgctaattc tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttgagcg aacgacctac accgaactga gatacctaca	4080
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt	4140
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccagggggaa acgcctggta	4200
tctttatagt cctgtcgggt ttcgccacct ctgacttgag cgtcgatttt tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ccttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgat	4440
ctatatcata atatgtacat ttatattggc tcatgtccaa tatgaccgcc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggc cattagttca tagcccatat	4560
atggagttcc gcgttacata acttacggta aatggccgc ctggctgacc gcccaacgac	4620
ccccgcccat tgacgtcaat aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtatttacgg taaactgcc acttggcagt acatcaagtg	4740
tatcatatgc caagtcgcc ccctattgac gtcaatgacg gtaaatggcc cgcttggcat	4800
tatgcccagt acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatgggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac	4980
caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc	5040

```

aaatgggcgg taggcgtgta cgggtgggagg tctatataag cagagctcgt ttagtgaacc 5100
gtcagatcac tagaagcttt attgcggttag tttatcacag ttaaattgct aacgcagtca 5160
gtgctttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc 5220
agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga 5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct 5340
attggtctta ctgacatcca ctttgccctt ctctccacag gtgtccactc ccagttcaat 5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga 5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgtccc 5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtccctga 5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg 5640
cagcttgaat gagaatatca ctgtcccaga caccgacgtt aattttctatg cctggaagag 5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc 5760
tgtcctgcgg ggccaggccc tgttggtcaa ctcttcccag ccgtgggagc ccctgcagct 5820
gcatgtggat aaagccgtca gtggccttcg cagcctcacc actctgcttc gggctctgcg 5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat 5940
cactgctgac actttccgca aactcttccg agtctactcc aatttcctcc ggggaaagct 6000
gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

```

```

<210> 211
<211> 6059
<212> DNA
<213> Artificial

```

```

<220>
<223> Description of Artificial Sequence: plasmid

```

```

<400> 211
ctagagtcga cccgggcggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60
ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120
atgtgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180
gttaacaaca acaattgcat tcattttatg tttcaggctc agggggagat gtgggaggtt 240
ttttaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300
ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360
gcgaatggac gcgccctgta gcggcgcatc aagcgcgggc ggtgtggtgg ttacgcgcag 420
cgtgaccgct acacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt 480
tctcgccacg ttgcgccggt ttccccgtca agctctaaat cgggggctcc ctttaggggt 540

```

ccgatttagt gctttacggc acctcgaccc caaaaaactt gattaggggtg atggttcacg 600  
 tagtggggcca tcgccctgat agacgggtttt tcgccctttg acgttggagt ccacgttctt 660  
 taatagtgga ctcttggtcc aaactggaac aacactcaac cctatctcgg tctattcttt 720  
 tgattttataa gggatttttg cgatttcggc ctattgggta aaaaatgagc tgatttaaca 780  
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtatttt 840  
 ctcccttacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900  
 ctgaaataac ctctgaaaga ggaacttggt taggtacctt ctgaggcgga aagaaccagc 960  
 tgtggaatgt gtgtcagtta ggggtgtggaa agtccccagg ctccccagca ggcagaagta 1020  
 tgcaaagcat gcatctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080  
 caggcagaag tatgcaaagc atgcatctca attagtcagc aaccatagtc ccgcccctaa 1140  
 ctccgcccctat ccgccccta actccgccc gttccgccc ttctccgccc catggctgac 1200  
 taattttttt tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260  
 agtgaggagg cttttttgga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320  
 cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcaggttc 1380  
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440  
 ctctgatgcc gccgtgttcc ggctgtcagc gcagggggcg ccggttcttt ttgtcaagac 1500  
 cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtggctggc 1560  
 caccaggggc gttccttgcg cagctgtgct cgacgttgct actgaagcgg gaagggactg 1620  
 gctgctattg ggcaagtgc cggggcagga tctcctgtca tctcaccttg ctctgcca 1680  
 gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg 1740  
 cccattcgac caccaagcga aacatcgcat cgagcgagca cgtactcgga tggaagccgg 1800  
 tcttgtcgat caggatgatc tggacgaaga gcatcagggg ctgcgcagcag ccgaactgtt 1860  
 cgccaggctc aaggcgcgca tgcccgacgg cgaggatctc gtctgaccc atggcgatgc 1920  
 ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg 1980  
 gctgggtgtg gcggaccgct atcaggacat agcgttggct acccgtgata ttgctgaaga 2040  
 gcttggcggc gaatgggctg accgcttctt cgtgctttac ggtatcgccg ctcccgattc 2100  
 gcagcgcac gccttctatc gccttcttga cgagttcttc tgagcggggac tctgggggtc 2160  
 gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt 2220  
 attttcatta catctgtgtg ttggtttttt gtgtgaatcg atagcgataa ggatccgcgt 2280  
 atggtgcact ctcagtacaa tctgctctga tgccgcatag ttaagccagc cccgacaccc 2340

gccaacacccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgctt atttttatag gttaatgtca tgataataat	2520
ggttttcttag acgtcagggtg gcacttttctg gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcattttgcc ttcctgtttt tgctcaccca gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gccccgaaga acgtttttcca atgatgagca cttttaaagt	2880
tctgctatgt ggcgcggtat tatcccgtat tgacgcgggg caagagcaac tcggtcgccg	2940
catacactat tctcagaatg acttggttga gtactcacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga tggaggcgga	3300
taaagttgca ggaccacttc tgcgctcggc ccttcgggtt ggctggttta ttgctgataa	3360
atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcaactgggc cagatggtaa	3420
gccctcccgt atcgtagtta tctacacgac ggggagtcag gcaactatgg atgaacgaaa	3480
tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt cagaccaagt	3540
ttactcatat atactttaga ttgatttaaa acttcatttt taatttaaaa ggatctaggt	3600
gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg	3660
agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt	3720
aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtgggtttgtt tgccggatca	3780
agagctacca actctttttc cgaaggtaac tggcttcagc agagcgcaga taccaaatac	3840
tgttcttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac	3900
atacctcgct ctgctaatac tgttaccagt ggctgctgcc agtggcgata agtcgtgtct	3960
taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcgg gctgaacggg	4020
gggttcgtgc acacagccca gcttggagcg aacgacctac accgaactga gatacctaca	4080
gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt	4140
aagcggcagg gtcggaacag gagagcgcac gagggagctt ccaggggggaa acgcctggta	4200

tctttatagt cctgtcgggt ttcgccacct ctgacttgag cgtcgatfff tgtgatgctc	4260
gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc	4320
cttttgctgg ccttttgctc acatggctcg acagatcttc aatattggcc attagccata	4380
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgat	4440
ctatatcata atatgtacat ttatatggc tcatgtccaa tatgaccgcc atgttggcat	4500
tgattattga ctagttatta atagtaatca attacggggc cattagttca tagcccatat	4560
atggagttcc gcgttacata acttacggta aatggccgcg ctggctgacc gcccacgac	4620
ccccgcccac tgacgtcaat aatgacgtat gttcccatag taacgccaat agggactttc	4680
cattgacgtc aatgggtgga gtatttacgg taaactgccc acttggcagt acatcaagtg	4740
tatcatatgc caagtccgcc ccctattgac gtcaatgacg gtaaattggc cgctggcat	4800
tatgccagc acatgacctt acgggacttt cctacttggc agtacatcta cgtattagtc	4860
atcgctatta ccatgggtgat gcggttttgg cagtacacca atgggcgtgg atagcggttt	4920
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac	4980
caaaatcaac gggactttcc aaaatgtcgt aacaactgcg atcgcccgcc ccgttgacgc	5040
aaatgggcgg taggcgtgta cgggtggagg tctatataag cagagctcgt ttagtgaacc	5100
gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca	5160
gtgcttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc	5220
agaagttggc cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga	5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct	5340
attggtctta ctgacatcca ctttgccctt ctctccacag gtgtccactc ccagttcaat	5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga	5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgtccc	5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcatc tgtgacagcc gagtccctga	5580
gaggtaacct ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg	5640
cagcttgaat gagaatatca ctgtcccaga caccaaagtt aatttctatg cctggaagag	5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc	5760
tgtcctgcgg ggccaggccc tgttggtcaa ctcttcccag ccgtgggagc ccctgcagct	5820
gcatgtggat aaagccgtcg agggccttcg cagcctcacc actctgcttc gggctctgcg	5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat	5940
cactgctgac actttccgca aactcttcg agtctactcc aatttcctcc ggggaaagct	6000

gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat 6059

<210> 212  
 <211> 6059  
 <212> DNA  
 <213> Artificial

<220>  
 <223> Description of Artificial Sequence: plasmid

<400> 212  
 ctagagtcga cccgggaggc cgcttccctt tagtgagggt taatgcttcg agcagacatg 60  
 ataagataca ttgatgagtt tggacaaacc acaactagaa tgcagtgaaa aaaatgcttt 120  
 atttgtgaaa tttgtgatgc tattgcttta tttgtaacca ttataagctg caataaacia 180  
 gttaacaaca acaattgcat tcattttatg tttcagggtc agggggagat gtgggaggtt 240  
 ttttaaagca agtaaaacct ctacaaatgt ggtaaaatcc gataaggatc gatccgggct 300  
 ggcgtaatag cgaagaggcc cgcaccgatc gcccttccca acagttgcgc agcctgaatg 360  
 gcgaatggac gcgccctgta ggcgcgcatt aagcgcggcg ggtgtggtgg ttacgcgcag 420  
 cgtgaccgct acacttgcca gcgccctagc gcccgctcct ttcgctttct tcccttcctt 480  
 tctcgccacg ttcgcgggct ttccccgtca agctctaaat cgggggctcc ctttaggggt 540  
 ccgatttagt gctttacggc acctcgacct caaaaaactt gattaggggtg atgggtcacg 600  
 tagtgggcca tcgccctgat agacggtttt tcgccctttg acgttgaggt ccacgttctt 660  
 taatagtgga ctcttggttc aaactggaac aacactcaac cctatctcgg tctattcttt 720  
 tgatttataa gggattttgc cgatttcggc ctattgggtt aaaaatgagc tgatttaaca 780  
 aaaatttaac gcgaatttta acaaaatatt aacgcttaca atttcctgat gcggtatttt 840  
 ctccttacgc atctgtgcgg tatttcacac cgcatacgcg gatctgcgca gcaccatggc 900  
 ctgaaataac ctctgaaaga ggaacttggg taggtacctt ctgaggcgga aagaaccagc 960  
 tgtggaatgt gtgtcagtta ggggtgtgga agtccccagg ctccccagca ggcagaagta 1020  
 tgcaaagcat gcatctcaat tagtcagcaa ccagggtgtg aaagtcccca ggctccccag 1080  
 caggcagaag tatgcaaagc atgcatctca attagtcagc aaccatagtc ccgcccctaa 1140  
 ctccgcccct cccgccccta actccgcccc gttccgcccc ttctccgccc catggctgac 1200  
 taattttttt tattttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt 1260  
 agtgaggagg ctttttttga ggcctaggct tttgcaaaaa gcttgattct tctgacacaa 1320  
 cagtctcgaa cttaaggcta gagccaccat gattgaacaa gatggattgc acgcagggtc 1380  
 tccggccgct tgggtggaga ggctattcgg ctatgactgg gcacaacaga caatcggctg 1440  
 ctctgatgcc gccgtgttcc ggctgtcagc gcaggggcgc ccggttcttt ttgtcaagac 1500

cgacctgtcc ggtgccctga atgaactgca ggacgaggca gcgcggctat cgtgggtggc	1560
cacgacgggc gttccttgcg cagctgtgct cgaogttgtc actgaagcgg gaagggactg	1620
gctgctattg ggcgaaagtgc cggggcagga tctcctgtca tctcaccttg ctccctgccga	1680
gaaagtatcc atcatggctg atgcaatgcg gcggctgcat acgcttgatc cggctacctg	1740
cccattcgac caccaagcga aacatcgcac cgagcgcagca cgtactcgga tggaaagccgg	1800
tcttgctgat caggatgatc tggacgaaga gcatcagggg ctgcgcgcag ccgaactggt	1860
cgccaggctc aaggcgcgca tgcccgcgcg cgaggatctc gtcgtgaccc atggcgatgc	1920
ctgcttgccg aatatcatgg tggaaaatgg ccgcttttct ggattcatcg actgtggccg	1980
gctgggtgtg gcggaccgct atcaggacat agcgttggt acccgtgata ttgctgaaga	2040
gcttggcggc gaatgggctg accgcttcct cgtgctttac ggtatcgccg ctcccgatcc	2100
gcagcgcac gccttctatc gccttcttga cgagttcttc tgagcgggac tctggggttc	2160
gaaatgaccg accaagcgac gcccaacctg ccatcacgat ggccgcaata aaatatcttt	2220
attttcatca catctgtgtg ttggtttttt gtgtgaatcg atagcgataa ggatccgcgt	2280
atgggtgcact ctcagtacaa tctgctctga tgccgcatag ttaagccagc cccgacaccc	2340
gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca	2400
agctgtgacc gtctccggga gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg	2460
cgcgagacga aagggcctcg tgatacgcct atttttatag gttaatgtca tgataataat	2520
ggtttcttag acgtcagggtg gcacttttctg gggaaatgtg cgcggaaccc ctatttgttt	2580
atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct gataaatgct	2640
tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg cccttattcc	2700
cttttttgcg gcattttgcc ttctgtttt tgctcaccca gaaacgctgg tgaaagtaaa	2760
agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc tcaacagcgg	2820
taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca cttttaaagt	2880
tctgctatgt ggcgcgggtat tatcccgat tgacgcggg caagagcaac tcggtcgccg	2940
catacactat tctcagaatg acttggttga gtactacca gtcacagaaa agcatcttac	3000
ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg ataacactgc	3060
ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt ttttgacaa	3120
catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg aagccatacc	3180
aaacgacgag cgtgacacca cgatgcctgt agcaatggca acaacgttgc gcaaactatt	3240
aactggcgaa ctacttactc tagcttccc gcaacaatta atagactgga tggaggcgga	3300

taaagttgca	ggaccacttc	tgcgctcggc	ccttccggct	ggctggttta	ttgctgataa	3360
atctggagcc	ggtgagcgtg	ggtctcgcgg	tatcattgca	gcactggggc	cagatggtaa	3420
gccctcccg	atcgtagtta	tctacacgac	ggggagtcag	gcaactatgg	atgaacgaaa	3480
tagacagatc	gctgagatag	gtgcctcact	gattaagcat	tggtaactgt	cagaccaagt	3540
ttactcatat	atactttaga	ttgatttaaa	acttcatttt	taatttaaaa	ggatctaggt	3600
gaagatcctt	tttgataatc	tcatgaccaa	aatcccttaa	cgtgagtttt	cgttccactg	3660
agcgtcagac	cccgtagaaa	agatcaaagg	atcttcttga	gatccttttt	ttctgcgcgt	3720
aatctgctgc	ttgcaaacaa	aaaaaccacc	gtaccagcgc	gtggtttggt	tgccggatca	3780
agagctacca	actctttttc	cgaaggtaac	tggcttcagc	agagcgcaga	taccaaatac	3840
tgttcttcta	gtgtagccgt	agttaggcca	ccacttcaag	aactctgtag	caccgcctac	3900
atacctcgct	ctgctaatac	tgttaccagt	ggctgctgcc	agtggcgata	agtcgtgtct	3960
taccgggttg	gactcaagac	gatagttacc	ggataaggcg	cagcggtcgg	gctgaacggg	4020
gggttcgtgc	acacagccca	gcttggagcg	aacgacctac	accgaactga	gatacctaca	4080
gcgtgagcta	tgagaaagcg	ccacgccttc	cgaagggaga	aaggcggaca	ggtatccggg	4140
aagcggcagg	gtcggaacag	gagagcgcac	gaggagcctt	ccagggggaa	acgcctggta	4200
tctttatagt	cctgtcgggt	ttcgccacct	ctgacttgag	cgtcgatttt	tgtgatgctc	4260
gtcagggggg	cggagcctat	ggaaaaacgc	cagcaacgcg	gcctttttac	ggttcctggc	4320
cttttgctgg	ccttttgctc	acatggctcg	acagatcttc	aatattggcc	attagccata	4380
ttattcattg	gttatatagc	ataaatcaat	attggctatt	ggccattgca	tacgttgat	4440
ctatatcata	atatgtacat	ttatattggc	tcatgtccaa	tatgaccgcc	atgttggcat	4500
tgattattga	ctagttatta	atagtaatca	attacggggg	cattagttca	tagcccatat	4560
atggagttcc	gcgttacata	acttacggta	aatggcccgc	ctggctgacc	gcccacgcac	4620
ccccgcccc	tgacgtcaat	aatgacgtat	gttcccatag	taacgccaat	agggactttc	4680
cattgacgtc	aatgggtgga	gtatttacgg	taaactgccc	acttggcagt	acatcaagtg	4740
tatcatatgc	caagtccgcc	ccctattgac	gtcaatgacg	gtaaatggcc	cgcctggcat	4800
tatgccaggt	acatgacctt	acgggacttt	cctacttggc	agtacatcta	cgtattagtc	4860
atcgctatta	ccatgggtgat	gcggtttttg	cagtacacca	atgggcgtgg	atagcggttt	4920
gactcacggg	gattttccaag	tctccacccc	attgacgtca	atgggagttt	gttttggcac	4980
caaaatcaac	gggactttcc	aaaatgtcgt	aacaactgcg	atcgcccgcc	ccgttgacgc	5040
aaatgggcgg	taggcgtgta	cgggtgggagg	tctatataag	cagagctcgt	ttagtgaacc	5100



gtcagatcac tagaagcttt attgcggtag tttatcacag ttaaattgct aacgcagtca	5160
gtgcttctga cacaacagtc tcgaacttaa gctgcagtga ctctcttaag gtagccttgc	5220
agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca ggtttaagga	5280
gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct	5340
attggctctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat	5400
tacagctctt aaggctagag tacttaatac gactcactat aggctagcct cgagcgcgga	5460
gatgggggtg cacgaatgtc ctgcctggct gtggcttctc ctgtccctgc tgtcgctccc	5520
tctgggcctc ccagtcctgg gcgccccacc acgcctcacc tgtgacagcc gagtccctgga	5580
gaggtacctc ttggaggcca aggaggccga gaatatcacg acgggctgtg ctgaacactg	5640
cagcttgaat gagaatatca ctgtcccaga caccgacgtt aatttctatg cctggaagag	5700
gatggaggtc gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcggaagc	5760
tgtcctgctg ggcagggccc tggtggtcaa ctcttcccag ccgtgggagc cctgcagct	5820
gcatgtggat aaagccgtcg agggccttcg cagcctcacc actctgcttc gggctctgctg	5880
agcccagaag gaagccatct cccctccaga tgcggcctca gctgctccac tccgaacaat	5940
cactgctgac actttccgca aactcttcctg agtctactcc aatttcctcc ggggaaagct	6000
gaagctgtac acaggggagg cctgcaggac aggggaccat catcaccatc accattgat	6059